

On compact hyperbolic Coxeter d -polytopes with $d + 4$ facets.

Anna Felikson¹, Pavel Tumarkin²

Abstract. We show that there is no compact hyperbolic Coxeter d -polytope with $d+4$ facets for $d \geq 8$. This bound is sharp: examples of such polytopes up to dimension 7 were found by Bugaenko [Bu1]. We also show that in dimension $d = 7$ the polytope with 11 facets is unique.

Contents

1 Preliminaries	2
2 Admissible Coxeter diagrams without small Lannér diagrams	9
3 Liftings	11
4 Absence of polytopes in dimensions ≥ 9	17
5 Absence of polytopes in dimension 8	20
6 Polytopes in dimension 7	26

Introduction

A Coxeter polytope in the spherical, hyperbolic or Euclidean space is a polytope whose dihedral angles are all integer submultiples of π . These polytopes are very important among the other acute-angled polytopes since a group generated by reflections with respect to the facets of a Coxeter polytope is discrete. On the other hand, a fundamental chamber of any discrete reflection group is a Coxeter polytope.

It is well known [C] that any spherical Coxeter polytope (containing no pair of opposite points of the sphere) is a simplex and any compact Euclidean Coxeter polytope is either a simplex or a direct product of simplices. See [C] for the description of these polytopes.

At the same time, hyperbolic Coxeter polytopes are still far from being classified. It is proved that no compact hyperbolic Coxeter polytope exists in dimensions $d \geq 30$ [V2], and no finite volume hyperbolic Coxeter polytope exists in dimensions $d \geq 996$ [P]. These bounds do not look sharp: examples of compact polytopes are known up to dimension 8 only [Bu1], [Bu2], and examples of finite volume polytopes are known up to dimension 21 only [V1], [VK], [Bo].

We will focus on compact Coxeter polytopes. Besides the restriction on the dimension and some series of examples [M], [ImH], [Al], [R], there exists a classification of compact hyperbolic Coxeter polytopes of certain combinatorial types. More precisely, simplices are classified in [L], d -polytopes with $d + 2$ facets were listed in [K] and [E2], and the classification of d -polytopes with $d + 3$ facets is contained in [T].

¹Partially supported by grants NS-5666.2006.1 and INTAS YSF-06-10000014-5916.

²Partially supported by grants MK-6290.2006.1, NS-5666.2006.1 and INTAS YSF-06-10000014-5766.

In this paper we examine a class of the next complexity, namely, compact hyperbolic Coxeter d -polytopes with $d + 4$ facets. We show that no such a polytope exists in hyperbolic space of dimension $d \geq 8$. We also stress that in dimensions $2 \leq d \leq 7$ these polytopes do exist [Bu1].

The paper is organized as follows. Section 1 is preparatory: we recall basic notions concerning Coxeter diagrams and combinatorics of simple polytopes. We recall also some facts connecting the combinatorial (metrical) properties of a face of a polytope to the combinatorial (metrical) properties of the polytope itself. Section 2 is devoted to Coxeter diagrams containing no Lannér diagram of order less than 5. In particular, we prove that a Coxeter diagram of any compact hyperbolic Coxeter polytope contains a Lannér subdiagram of order less than 5. In Section 3 we develop a theory of liftings to connect the combinatorics of a face of a Coxeter polytope to a subdiagram of the Coxeter diagram of this polytope. Finally, in Sections 4 and 5 we use all these tools to show the absence of the polytopes in dimensions $d \geq 8$.

The paper was partially written in the Max Plank Institute for Mathematics in Bonn. The authors are grateful to the Institute for hospitality.

1 Preliminaries

In this section we list the essential facts about Coxeter diagrams, Gale diagrams and diagrams of missing faces. Concerning Coxeter diagrams we follow mainly [V3] and [V4]. For the details about Gale diagrams see [G]. See also [E1] for an overview about Coxeter polytopes, Gale diagrams and diagrams of missing faces. At the end of the section we recall a recent result of Allcock [Al] which states that Coxeter polytopes often have some Coxeter faces and describes the Coxeter diagrams of that faces.

Coxeter diagrams

1. An *abstract Coxeter diagram* Σ is a finite 1-dimensional simplicial complex with weighted edges, where weights w_{ij} are positive, and if $w_{ij} < 1$ then $w_{ij} = \cos \frac{\pi}{m_{ij}}$ for some integer $m_{ij} \geq 3$. A *subdiagram* of Σ is a subcomplex with the same weights as in Σ . The *order* $|\Sigma|$ of the diagram Σ is the number of nodes of Σ .

If Σ_1 and Σ_2 are subdiagrams of an abstract Coxeter diagram Σ , we denote by $\langle \Sigma_1, \Sigma_2 \rangle$ a subdiagram of Σ spanned by all nodes of Σ_1 and Σ_2 . Denote by $\Sigma_1 \setminus v$ and $\Sigma_1 \setminus \Sigma_2$ the subdiagrams of the diagram Σ_1 spanned by all nodes of Σ_1 except either v or nodes of Σ_2 respectively.

Given an abstract Coxeter diagram Σ with nodes $v_1, \dots, v_n$ and weights w_{ij} , we construct a symmetric $n \times n$ matrix $G(\Sigma) = (g_{ij})$, where $g_{ii} = 1$, $g_{ij} = -w_{ij}$ if v_i and v_j are adjacent, and $g_{ij} = 0$ otherwise. By $\det(\Sigma)$ and a signature of Σ we mean the determinant and the signature of $G(\Sigma)$.

We can draw edges of Coxeter diagram in the following way: if the weight w_{ij} equals $\cos(\frac{\pi}{m_{ij}})$, v_i and v_j are joined by an $(m_{ij} - 2)$ -fold edge or a simple edge labeled by m_{ij} ; if $w_{ij} = 1$, v_i and v_j are joined by a bold edge; if $w_{ij} > 1$, v_i and v_j are joined by a dotted edge labeled by w_{ij} (or without any label).

We write $[v_i, v_j] = m_{ij}$ if $w_{ij} = \cos(\frac{\pi}{m_{ij}})$, and $[v_i, v_j] = \infty$ if $v_i v_j$ is a dotted edge. We write $[v_i, v_j] = 2$ if v_i and v_j are not joined.

An abstract Coxeter diagram Σ is *elliptic* if $G(\Sigma)$ is positive definite; Σ is *parabolic* if any indecomposable component of $G(\Sigma)$ is degenerate and positive semidefinite; a connected diagram Σ is a *Lannér* diagram if Σ is neither elliptic nor parabolic but any proper subdiagram of Σ is elliptic; Σ is *hyperbolic* if Σ is indefinite with negative inertia index equal to 1; Σ is

superhyperbolic if its negative inertia index is greater than 1; Σ is *admissible* if Σ contains no parabolic subdiagrams and Σ is not superhyperbolic.

Table 1 contains the list of elliptic and connected parabolic Coxeter diagrams with their standard notation. Lannér diagrams are listed in [V4, Table 3]. In particular, there are finitely many Lannér diagrams of order greater than 3, and the maximal order of a Lannér diagram is 5. We represent the list of Lannér diagrams of order 4 and 5 in Table 2.

2. It is convenient to describe Coxeter polytopes by their Coxeter diagrams. Let P be a Coxeter polytope with facets $f_1, \dots, f_r$. The Coxeter diagram $\Sigma(P)$ of the polytope P is a diagram with nodes $v_1, \dots, v_r$; two nodes v_i and v_j are not joined if f_i is orthogonal to f_j ; v_i and v_j are joined by an edge with weight

$$w_{ij} = \begin{cases} \cos \frac{\pi}{k}, & \text{if } f_i \text{ and } f_j \text{ form a dihedral angle } \frac{\pi}{k}; \\ 1, & \text{if } f_i \text{ is parallel to } f_j; \\ \cosh \rho, & \text{if } f_i \text{ and } f_j \text{ diverge and } \rho \text{ is the distance from } f_i \text{ to } f_j. \end{cases}$$

If $\Sigma = \Sigma(P)$ then $M(\Sigma)$ coincides with the Gram matrix of unit normal vectors to the facets of P .

It is shown in [V3] that if $\Sigma = \Sigma(P)$ is a Coxeter diagram of a compact d -dimensional hyperbolic polytope P , then Σ is an admissible connected hyperbolic diagram with positive inertia index equal to d . In particular, Σ contains no bold edges and other parabolic subdiagrams. Elliptic subdiagrams of Σ are in one-to-one correspondence with faces of P : a k -face F corresponds to an elliptic subdiagram Σ_F of order $d - k$ whose nodes correspond to the $(d - 1)$ -faces containing F .

3. Given a Coxeter diagram Σ , it is easy to check if Σ is superhyperbolic or not. However, this requires the computation of a signature of a rather big matrix if the order of Σ is not small. In the case when Σ is a union of two subdiagrams either joined by only one edge or having a unique node in common, there exists a more effective way to state that Σ is superhyperbolic [V2].

Let T be a subdiagram of Σ such that $\det(\Sigma \setminus T) \neq 0$. A *local determinant* of Σ on a subdiagram T is $\det(\Sigma, T) = \frac{\det(\Sigma)}{\det(\Sigma \setminus T)}$.

Proposition 1.1 ([V2], Prop. 12). *If a Coxeter diagram Σ consists of two subdiagrams Σ_1 and Σ_2 having a unique vertex v in common, and nodes of $\Sigma_1 \setminus v$ do not attach to $\Sigma_2 \setminus v$, then*

$$\det(\Sigma, v) = \det(\Sigma_1, v) + \det(\Sigma_2, v) - 1.$$

Proposition 1.2 ([V2], Prop. 13). *If a Coxeter diagram Σ is generated by disjoint subdiagrams Σ_1 and Σ_2 joined by a single edge v_1v_2 , then*

$$\det(\Sigma, \langle v_1, v_2 \rangle) = \det(\Sigma_1, v_1) \det(\Sigma_2, v_2) - w_{12}^2,$$













where w_{12} is the weight of the edge v_1v_2 .

Proposition 1.3 ([V2], Prop. 15). *Suppose that a Coxeter diagram Σ is a union of two disjoint hyperbolic subdiagrams Σ_1 and Σ_2 joined by a unique edge v_1v_2 and that the subdiagrams $\Sigma_1 \setminus v_1$ and $\Sigma_2 \setminus v_2$ are elliptic. Assume that in addition one of the following conditions holds:*

- 1) v_1v_2 is a simple edge and $\det(\Sigma_1, v_1) \det(\Sigma_2, v_2) > \frac{1}{4}$;
- 2) v_1v_2 is a double edge and $\det(\Sigma_1, v_1) \det(\Sigma_2, v_2) > \frac{1}{2}$.

Then the diagram Σ is superhyperbolic.

Table 1: Connected elliptic and parabolic Coxeter diagrams are listed in the left and in the right column respectively

A_n ($n \geq 1$) 	$\tilde{A}_1$  $\tilde{A}_n$ ($n \geq 2$) 
$B_n = C_n$ ($n \geq 2$) 	$\tilde{B}_n$ ($n \geq 3$)  $\tilde{C}_n$ ($n \geq 2$) 
D_n ($n \geq 4$) 	$\tilde{D}_n$ ($n \geq 4$) 
$G_2^{(m)}$ 	$\tilde{G}_2$ 
F_4 	$\tilde{F}_4$ 
E_6 	$\tilde{E}_6$ 
E_7 	$\tilde{E}_7$ 
E_8 	$\tilde{E}_8$ 
H_3 	
H_4 	

In [V2, Table 2] Vinberg listed some useful local determinants. We use Propositions 1.1– 1.3 together with Table 2 of [V2] throughout this paper not referring to them every time. We also use the fact that all local determinants of Lannér diagrams shown in Table 2 (of the present paper) on their open vertices (see the definition below) are greater than 0.95, this fact can be checked by a direct computation. In particular, we will use the following corollary of Prop. 1.3:

Proposition 1.4. *Suppose that a diagram Σ consists of two disjoint Lannér diagrams L_1 and L_2 of order 5 each joined by a unique non-dotted edge v_1v_2 , such that $L_i \setminus v_i$ (where $v_i \in L_i$, $i = 1, 2$) is of the type H_4 or F_4 . Then Σ is superhyperbolic.*


4. Let Σ be an abstract Coxeter diagram and v be a node of Σ . Suppose that it is possible to construct an abstract Coxeter diagram $\langle \Sigma, x \rangle$, $x \notin \Sigma$, such that x is joined with v in $\langle \Sigma, x \rangle$ and for any non-elliptic subdiagram $\Sigma' \subset \langle \Sigma, x \rangle$ the intersection $\Sigma' \cap \Sigma = \Sigma' \setminus x$ is non-elliptic, too. Then v is called an *open vertex* of Σ .

The vertex v is called *doubly open* if it is possible to construct a connected Coxeter diagram $\langle \Sigma, x_1 \rangle$, $x_1 \notin \Sigma$, such that x_1 is joined with v in $\langle \Sigma, x_1 \rangle$ and x_1 is an open vertex of $\langle \Sigma, x_1 \rangle$.

If $v \in \text{Sigma}$ is not an open vertex of a Lannér diagram Σ and $x \notin \Sigma$ is a node joined with v , then the diagram $\langle \Sigma, x \rangle$ contains a non-elliptic subdiagram M such that $x, v \in M$. Indeed, if x is not connected with $\Sigma \setminus v$, the statement is obvious. If we add some edges connecting x and $\Sigma \setminus v$ then the non-elliptic subdiagrams of $\langle \Sigma, x \rangle$ remain non-elliptic.

In [E1, Tabelle 2] Esselmann listed all Lannér diagrams of order 4 and 5 containing open vertices. In Table 2 we list Lannér diagrams of order 4 and 5 and introduce the notation. The open vertices are marked black, the doubly open vertices are encircled.

Table 2: Lannér diagrams of order 4 and 5. Open vertices are marked black, doubly open vertices are marked by additional circles. The upper index equals the order of the diagram.



Proposition 1.5. *Any Lannér diagram of order 4 contains at least two subdiagrams of the types H_3 or B_3 .*

This follows directly from the classification of Lannér diagrams.

Proposition 1.6 ([V2], Prop. 2). *Any two Lannér diagrams of Σ are joined by at least one edge.*

The statement is obvious: if it is false then Σ is a superhyperbolic diagram.

Gale diagrams and diagrams of missing faces

1. As it is shown in [V3, cor. of Th. 3.1], any compact hyperbolic Coxeter polytope is simple (i.e. a k -dimensional face of a d -polytope is an intersection of exactly $d - k$ facets). In this section we discuss the combinatorics of simple polytopes. We abuse notation by writing “ d -polytope” instead of “ d -dimensional polytope”.

Every combinatorial type of simple d -polytope with $d + k$ facets can be represented by its *Gale diagram*. This consists of $d + k$ points $a_1, \dots, a_{d+k}$ of $(k - 2)$ -dimensional unit sphere $\mathbb{S}^{k-2} \subset \mathbb{R}^{k-1}$ centered at the origin. Each point a_i corresponds to a facet f_i of P . The combinatorial type of a simple convex polytope can be read off from the Gale diagram in the following way: for any $J \subset \{1, \dots, d + k\}$ the intersection of facets $\{f_j | j \in J\}$ is a face of P if and only if the origin is contained in the interior of $\text{conv}\{a_j | j \notin J\}$ (where $\text{conv } X$ is a convex hull of the set X).

The set of points $\{a_1, \dots, a_{d+k}\} \subset \mathbb{S}^{k-2}$ is a Gale diagram of some d -dimensional polytope P with $d + k$ facets if and only if every open halfspace H^+ in $\mathbb{R}^{k-1}$ bounded by a hyperplane through the origin contains at least two of points $a_1, \dots, a_{d+k}$.

Two Gale diagrams are called *isomorphic* if the corresponding polytopes are combinatorially equivalent.

Let P be a simple polytope. The facets $f_1, \dots, f_m$ of P compose a *missing face* of P if $\bigcap_{i=1}^m f_i = \emptyset$ but any proper subset of $\{f_1, \dots, f_m\}$ has a non-empty intersection. Clearly, if a set of facets have no common intersection then this set of facets contains at least one missing face. This leads to the following lemma:

Lemma 1.1. *Let P be a simple d -polytope with $d + k$ facets and let $\{a_1, \dots, a_{d+k}\} \subset \mathbb{S}^{k-2}$ be a Gale diagram of P . Let H^+ be any open halfspace bounded by a hyperplane through the origin. Then H^+ contains a set $I \subset \{a_1, \dots, a_{d+k}\}$ which corresponds to a missing face of P .*

2. If $k = 2$ then the Gale diagram of P is one-dimensional, i.e. vertices a_i of the diagram lie on the 0-dimensional unit sphere. In other words, each of the points a_i coincides either with the point -1 or with the point 1 . This leads to the following

Proposition 1.7 ([G]). *A simple d -polytope with $d + 2$ facets is a direct product of two simplices $\Delta^{n-k} \times \Delta^k$ (where $0 \leq k \leq \lfloor n/2 \rfloor$ and Δ^m stays for an m -dimensional simplex).*

As it is shown in [E2], any compact Coxeter d -polytope with $d + 2$ facets is either a simplicial prism or a product of two triangles. Coxeter prisms are listed in [K], and the remaining Coxeter polytopes of this type are listed in [E2] (there are 7 of them). We call these 7 polytopes *Esselmann polytopes*.

If $k = 3$ then the Gale diagram of P is two-dimensional, i.e. vertices a_i of the diagram lie on the unit circle.

A *standard Gale diagram* of simple d -polytope with $d + 3$ facets consists of vertices $v_1, \dots, v_m$ of regular m -gon (m is odd) in $\mathbb{R}^2$ centered at the origin, which are labeled according to the following rules:

- 1) Each label is a positive integer, the sum of labels equals $d + 3$.
- 2) The vertices that lie in any open halfspace bounded by a line through the origin have labels whose sum is at least two.

It is easy to check (see, for example, [G]) that any two-dimensional Gale diagram is isomorphic to some standard diagram. Two simple d -polytopes with $d + 3$ facets are combinatorially equivalent if and only if their standard Gale diagrams are congruent.

For $k > 3$ there is no notion of standard Gale diagram known to us. We will use another diagram to encode the combinatorics of a simple polytope.

A *diagram of missing faces* is a finite set D with a specified collection $\mathcal{M}_D$ of subsets of D satisfying the following condition: if $M, M' \in \mathcal{M}_D$ then $M' \not\subset M$. The order $|M|$ of M is the number of elements in M . Elements of $\mathcal{M}_D$ are called *missing faces* of D .

A diagram $D_1 \subset D$ of missing faces is a *subdiagram* of D if for any $M \subset D_1$ the following holds: $M \in \mathcal{M}_{D_1}$ if and only if $M \in \mathcal{M}_D$.

To represent a diagram of missing faces we draw a point (vertex) for each element of D and encircle a set of points corresponding to M if and only if $M \in \mathcal{M}_D$.

3. For any simple polytope P we construct a diagram of missing faces $D(P)$ in the following way: the elements correspond to the facets of P , a set of elements is a missing face of $D(P)$ if and only if the corresponding facets compose a missing face of P .

The combinatorics of P can be recovered from $D(P)$: the set of facets have a non-empty intersection if and only if the corresponding set in $D(P)$ contains no missing faces.

Lemma 1.2. *Let P be a simple polytope and $D(P)$ be a diagram of missing faces of P . For any $M \in \mathcal{M}_{D(P)}$ there exists $M' \in \mathcal{M}_{D(P)}$ such that $M \cap M' = \emptyset$.*

Proof. Consider a Gale diagram $G(P)$ of P . It consists of several points on a sphere $\mathbb{S}^d$ for some d . Let $\overline{M}$ be the points of $G(P)$ corresponding to the vertices of M . Since M is a missing face, $\text{conv}(G(P) \setminus \overline{M})$ does not contain the origin. Equivalently, there exists a hyperplane H through 0 separating a halfspace H^+ containing $G(P) \setminus \overline{M}$. Denote by H^- another halfspace with respect to H . By Lemma 1.1, H^- contains at least one subset corresponding to some missing face. Since this subset belongs to $\overline{M}$, and no missing face contains another missing face, we obtain that H separates $G(P) \setminus \overline{M}$ from $\overline{M}$, i.e. $H^- \cap G(P) = \overline{M}$. By Lemma 1.1, H^+ contains at least one subset corresponding to some missing face M' . Clearly, M and M' are disjoint and the lemma is proved. $\square$

We also need the following two Propositions:

Proposition 1.8 ([E1], Lemma 1.6). *Let P be a simple polytope and f be a facet of P . Let $\{f_1, \dots, f_k\}$ be the set of all facets of P such that $f_i \cap f \neq \emptyset$ and $f_i \neq f$. Denote $f'_i = f_i \cap f$ for $i = 1, \dots, k$, and for any subset $F \subseteq \{f_1, \dots, f_k\}$ denote $F' = \{f'_i \mid f_i \in F\}$.*

The set of faces F' is a missing face of f if and only if either

- 1) $\{f\} \cup F$ is a missing face of P or
- 2) F is a missing face of P and F contains no proper subset F_0 such that $\{f\} \cup F_0$ is a missing face of P .

Proposition 1.9 ([E1], Lemma 1.9). *Let P be a simple polytope, and let f be any facet of P . There exists a missing face of P containing f .*

Lannér diagrams and missing faces

Let P be a compact Coxeter polytope in $\mathbb{H}^n$ and $\Sigma(P)$ be the Coxeter diagram of P . Let L be any Lannér subdiagram of $\Sigma(P)$. By definition of Lannér diagrams, the facets corresponding to L compose a missing face of P (and any missing face of P corresponds to some Lannér diagram in $\Sigma(P)$). Thus, a diagram of missing faces $D(P)$ can be easily reconstructed by $\Sigma(P)$: to obtain $D(P)$ one should take $\Sigma(P)$, encircle all Lannér diagrams and delete all edges.

In the same way we can construct a diagram $D(\Sigma)$ of missing faces for any admissible Coxeter diagram Σ .

The correspondence “Lannér diagrams $\longleftrightarrow$ missing faces” shows in particular that if P is a compact hyperbolic Coxeter polytope, then P has no missing faces of order greater than 5.

Faces of Coxeter polytopes

Let P be a compact hyperbolic Coxeter d -polytope, and denote by Σ its Coxeter diagram. Let S_0 be an elliptic subdiagram of Σ . By [V3, Th. 3.1], S_0 corresponds to a face of P of dimension $d - |S_0|$. Denote this face by $P(S_0)$. $P(S_0)$ itself is an acute-angled polytope [An], but it may not be a Coxeter polytope. Borchers obtained the following sufficient condition for $P(S_0)$ to be a Coxeter polytope.

Proposition 1.10 ([Bo], Example 5.6). *Suppose P is a Coxeter polytope with diagram Σ , and $S_0 \subset \Sigma$ is an elliptic subdiagram that has no A_n or D_5 component. Then $P(S_0)$ itself is a Coxeter polytope.*

Facets of $P(S_0)$ correspond to those nodes of Σ that compose an elliptic diagram together with S_0 . The following result of Allcock shows how to compute dihedral angles of $P(S_0)$.

We say that a node of Σ *attaches* to S_0 if it is joined with any node of S_0 by an edge of any type. Let a and b be the facets of $P(S_0)$ coming from facets A and B of P , i.e. $a = A \cap P(S_0)$ and $b = B \cap P(S_0)$. Denote by v_A and v_B the nodes of Σ corresponding to the facets A and B . Then the angles of $P(S_0)$ can be computed in the following way.

Proposition 1.11 ([Al], Th. 2.2.). *Under the hypotheses of Prop. 1.10,*

- (1) *If neither v_A nor v_B attaches to S_0 , then $\angle ab = \angle AB$.*
- (2) *If just one of v_A and v_B attaches to S_0 , say to the component S_0^i , then*
 - (a) *if $A \perp B$ then $a \perp b$;*
 - (b) *if v_A and v_B are joined and adjoining v_A and v_B to S_0^i yields a diagram B_k (resp. D_k , E_8 or H_4) then $\angle ab = \pi/4$ (resp. $\pi/4$, $\pi/6$ or $\pi/10$);*
 - (c) *otherwise, a and b do not meet.*
- (3) *If v_A and v_B attach to different components of S_0 , then*
 - (a) *if $A \perp B$ then $a \perp b$;*
 - (b) *otherwise, a and b do not meet.*
- (4) *If v_A and v_B attach to the same component of S_0 , say S_0^i , then*
 - (a) *if v_A and v_B are not joined and $S_0^i \cup \{A, B\}$ is a diagram E_6 (resp. E_8 or F_4) then $\angle ab = \pi/3$ (resp. $\pi/4$ or $\pi/4$);*
 - (b) *otherwise, a and b do not meet.*

Let $w \in \Sigma$ be a *neighbor* of S_0 , so that w attaches to S_0 by some edges. We call w *good* if $\langle S_0, w \rangle$ is an elliptic diagram, and *bad* otherwise. We denote by $\overline{S_0}$ the subdiagram of Σ consisting of nodes corresponding to facets of $P(S_0)$. The diagram $\overline{S_0}$ is spanned by good neighbors of S_0 and by all nodes not joined to S_0 . If $P(S_0)$ is a Coxeter polytope, denote its Coxeter diagram by Σ_{S_0} . By ordinary edge we mean non-dotted edges. By simple edges we mean 1-fold edges. By empty edge we mean two nodes which are not joined.

For any node $v \in \Sigma$ which is not a bad neighbor of S_0 (i.e. $v \in \overline{S_0}$) we denote by $\tilde{v}$ the corresponding node of Σ_{S_0} .

If Σ_{S_0} does not differ from $\overline{S_0}$, we consider the diagram Σ_{S_0} as a subdiagram of Σ , and we do not differ v and $\tilde{v}$.

Corollary 1.1. *Under the hypotheses of Prop. 1.10,*

- (a) *If S_0 is of the type H_4 , F_4 , or $G_2^{(m)}$ for $m \geq 6$, or any other diagram having no good neighbors, then $\overline{S}_0 = \Sigma_{S_0}$.*
- (b) *If S_0 is of the type H_3 , then $\overline{S}_0$ may be obtained from Σ_{S_0} by replacing some dotted edges by ordinary edges.*
- (c) *If S_0 is of the type $G_2^{(5)}$, then $\overline{S}_0$ may be obtained from Σ_{S_0} by replacing some edges labeled by 10 by simple edges, and some dotted edges by ordinary edges.*
- (d) *If S_0 is of the type B_n , $n \geq 3$, then $\overline{S}_0$ may be obtained from Σ_{S_0} by replacing some double edges by simple edges, and some dotted edges by ordinary edges.*
- (e) *If S_0 is of the type $B_2 = G_2^{(4)}$, then $\overline{S}_0$ may be obtained from Σ_{S_0} by replacing some double edges by simple edges, and some dotted edges by ordinary or empty edges.*

The corollary immediately follows from Prop. 1.11. One should only notice that all neighbors of diagrams listed in item (a) are bad.

Corollary 1.2. *Under the hypotheses of Prop. 1.10, let $S_1 \subset \Sigma_{S_0}$ be a subdiagram of the type $G_2^{(m)}$, where $m \neq 4, 10$, and let S'_1 be the corresponding subdiagram of $\overline{S}_0$. Then no node of $S'_1 \subset \Sigma$ is a good neighbor of S_0 .*

In particular, any subdiagram $S_2 \subset \Sigma_{S_0}$ of the type F_4 , H_4 , H_3 or $G_2^{(m)}$, where $m \neq 4, 10$, corresponds to a subdiagram of the same type in $\overline{S}_0$.

Any Lannér subdiagram $L \subset \Sigma_{S_0}$ of order 5 corresponds to a Lannér subdiagram of $\overline{S}_0$.

Lemma 1.3. *Suppose that $S_0 \subset \Sigma$ is an elliptic subdiagram and $|S_0| < d$. Then S_0 has at most $|\Sigma| - d - 1$ bad neighbors. In particular, if P has $d + 4$ facets, then any elliptic subdiagram of Σ of order less than d has at most 3 bad neighbors.*

Proof. The lemma follows from the fact that a d -polytope has at least $d + 1$ facets. □

Lemma 1.4. *Let $S \subset \Sigma$ be an elliptic subdiagram containing no component of the types A_n and D_5 , and let v be a bad neighbor of S . Then v is joined with each Lannér subdiagram of $\overline{S}$.*

The statement immediately follows from Proposition 1.6.

2 Admissible Coxeter diagrams without small Lannér diagrams

A Lannér diagram (a missing face) L is *small* if $|L| < 5$. A missing face M is *large* if $|M| > 5$.

Lemma 2.1. *Let Σ be a connected admissible Coxeter diagram containing no small Lannér subdiagrams. Suppose that each node of Σ belongs to some Lannér subdiagram of Σ . Then $|\Sigma| \leq 10$. If $|\Sigma| = 10$ then Σ is one of the three diagrams shown in Fig. 1.*

If in addition $\det(\Sigma) = 0$ then $\Sigma = \Theta_1$ (see Fig. 1).

Proof. Since Σ is a connected diagram and Σ contains no small Lannér subdiagrams, any edge of Σ is either a simple edge, or a double edge, or a triple edge. Consider two cases: either Σ contains $\mathcal{L}_5^5$ or it does not (see Table 2 for the notation).




Figure 1: Diagrams without small Lannér subdiagrams.

Case 1. Suppose that Σ contains $\mathcal{L}_5^5$. Clearly, only the open vertex of $\mathcal{L}_5^5$ can be joined with some other nodes of Σ , otherwise Σ contains a parabolic subdiagram $\tilde{B}_4$, or $\tilde{B}_5$, or a small Lannér diagram. By the similar reason, the diagram $\Sigma \setminus \mathcal{L}_5^5$ is a linear diagram without double edges. Since any node of Σ belongs to some Lannér diagram, for $|\Sigma| > 9$ such a diagram is superhyperbolic.

Case 2. Suppose that Σ does not contain $\mathcal{L}_5^5$.

- Any triple edge of Σ is incident to one of the leaves of Σ . The other vertex of the triple edge has valency two.

Proof: the statement follows immediately from the absence of small Lannér subdiagrams.

- Σ contains no double edge.

Proof: Suppose that v_1v_2 is a double edge of Σ . Assume that v_1 is a leaf of Σ . By Prop. 1.9, the node v_1 belongs to some Lannér diagram L . Clearly, L contains v_2 and L is a linear Lannér diagram. Let v be any node of $\Sigma \setminus L$ connected to L . Then $\langle L, v \rangle$ contains either parabolic or small Lannér subdiagram.

Therefore, neither v_1 nor v_2 is a leaf of Σ . Then Σ contains edges v_0v_1 and v_2v_3 ($v_0 \neq v_3$, otherwise $\langle v_1, v_2, v_3 \rangle$ contains either a Lannér or a parabolic subdiagram). Since Σ does not contain neither small Lannér diagrams nor parabolic diagrams, these edges are simple. Hence, $\langle v_0, v_1, v_2, v_3 \rangle$ is a diagram of the type F_4 . If v_4 is any node of Σ connected to $\langle v_0, v_1, v_2, v_3 \rangle$, then $\langle v_0, v_1, v_2, v_3, v_4 \rangle$ either is a diagram of the type $\mathcal{L}_5^5$ or it contains a small Lannér or a parabolic subdiagram, which contradicts the assumption.

- Σ is a tree.

Proof: By the previous two statements, any minimal cycle in Σ is a parabolic diagram $\tilde{A}_n$.

Remove from Σ all leaves that belong to triple edges. Denote the obtained diagram by Σ' . It follows from above that Σ' is a simply laced (i.e. it contained simple edges only) tree. It follows from the classification of parabolic diagrams that a simply laced tree without parabolic subdiagrams is an elliptic diagram. Therefore, Σ' is one of A_n, D_n, E_6, E_7, E_8 .

Now, append triple edges to some of the leaves of Σ' . If $\Sigma' = A_n$ or D_{n+1} for $n \geq 9$, then some vertices of Σ belong to no Lannér diagram. For the remaining cases we have either $|\Sigma| < 10$, or Σ is one of the diagrams listed in the lemma, or Σ is superhyperbolic. A direct calculation of determinants shows that $\det \Sigma = 0$ if and only if $\Sigma = \Theta_1$. □

Corollary 2.1. *Let P be a compact Coxeter polytope. Then $\Sigma(P)$ contains a small Lannér diagram.*

Proof. Suppose that $\Sigma(P)$ contains no small Lannér diagrams. Then, by Lemma 1.2, $\Sigma(P)$ contains two disjoint Lannér diagrams L_1 and L_2 of order 5, and hence, $|\Sigma(P)| \geq 10$. A subdiagram $\langle L_1, L_2 \rangle$ is connected, otherwise it is superhyperbolic. It follows from Lemma 2.1 that $\Sigma(P)$ is one of the diagrams listed in Fig. 1. In all three cases $\Sigma(P)$ contains a Lannér subdiagram which intersects any other Lannér subdiagram of $\Sigma(P)$. By Lemma 1.2, $\Sigma(P)$ cannot be a Coxeter diagram of a compact Coxeter polytope and the statement is proved. $\square$

3 Liftings

Let D be a diagram of missing faces and Σ be an admissible Coxeter diagram. Σ is a *0-lifting* of D if there is a bijection $\phi : D \rightarrow \Sigma$ such that $M \in \mathcal{M}_D$ if and only if $\phi(M)$ is a Lannér diagram. ϕ is called a *lifting bijection*.

Σ is a *k-lifting* of D ($k \in \mathbb{N}$) if Σ contains a subset A of additional nodes satisfying

- 1) $|A| = k$;
- 2) There exists an injection $\phi : D \rightarrow \Sigma$ taking bijectively D to $\Sigma \setminus A$. ϕ is called a *lifting injection*;
- 3) For any Lannér diagram L of Σ the set $\phi^{-1}(L \setminus A)$ contains a missing face of D ;
- 4) For any $M \in \mathcal{M}_D$ there exists a Lannér diagram $L \subset \langle \phi(M), A \rangle$ containing $\phi(M)$.
- 5) For any set $\{a_1, \dots, a_r\} \subset A$ the diagram $\Sigma \setminus \{a_1, \dots, a_r\}$ is not a $(k - r)$ -lifting of D .

In [E1] a 0-lifting of D is called a “hyperbolic realization” of D . A 0-lifting satisfies the general definition of a k -lifting. When the number k is not important, we write “lifting” instead of “ k -lifting”.

Let Σ be a lifting of D . Suppose that L is a Lannér diagram of Σ and for any $M \in \mathcal{M}_D$ we have $L \not\subset \langle \phi(M), A \rangle$. We say that L is an *additional* Lannér diagram of the lifting Σ .

Let D be an abstract diagram of missing faces, Σ be an abstract Coxeter diagram and ϕ be an injection $\phi : D \rightarrow \Sigma$. Suppose that there exists $\Sigma' \subset \Sigma$ which is a non-elliptic subdiagram not containing $\phi(M)$ for any $M \in \mathcal{M}_D$. Then we say that Σ' is a *conflicting* subdiagram (and conclude that Σ is not a lifting of D with given injection).

Notice that $\phi(M)$ is a subset of the set of nodes of Σ . When we mean a Coxeter diagram spanned by the nodes of $\phi(M)$, we write $\langle \phi(M) \rangle$ (compare the notation introduced in Section 1).

Lemma 3.1. *Let D be a diagram of missing faces and $D_1 \subset D$ be a subdiagram. Let Σ be a k -lifting of D . Then Σ contains a subdiagram Σ_1 which is a k_1 -lifting of D_1 for some $k_1 \leq k$.*

Proof. Let A be the set of additional nodes of Σ . Take $\phi|_{D_1}$ for the lifting injection of D_1 . By definition of liftings, $\langle \phi(D_1), A \rangle$ contains some k_1 -lifting of D_1 , and we have $k_1 \leq k$. $\square$

The proof of the following lemma is based on the multiple usage of Proposition 1.8.

Lemma 3.2. *Let $P \subset \mathbb{H}^d$ be a simple Coxeter polytope and let f be an m -face of P . Let $D(f)$ be a diagram of missing faces of f . Then $\Sigma(P)$ contains a subdiagram Σ_0 which is a k -lifting of $D(f)$ for some $k \leq d - m$.*

Proof. Let $f_1, \dots, f_{d-m}$ be the facets of P containing f . Let $F_1 = f_1$ and define inductively $F_i = F_{i-1} \cap f_i$. Clearly, F_i is a facet of F_{i-1} and $F_{d-m} = f$.

If $m = d$, the lemma is trivial. Assume by induction that the statement holds for any face of dimension greater than m . Let us prove for the m -face f . By the induction assumption, $\Sigma(P)$ contains a subdiagram Σ_{r-1} which is a k -lifting of $D(F_{d-m-1})$, where $k \leq d - (m + 1)$. The following claim completes the proof.

Claim: either Σ_{r-1} contains a k -lifting of f or $\langle \Sigma_{r-1}, v_{d-m} \rangle$ contains a $(k + 1)$ -lifting of f , where v_{d-m} is a node of $\Sigma(P)$ corresponding to the facet f_{d-m} .

Proof of the claim:

Let $\Pi_1, \dots, \Pi_s$ be the facets of F_{d-m-1} . Denote by J a set of indices such that $\Pi_i \cap f_{d-m} \neq \emptyset$ for $i \in J$, and denote $\pi_i = \Pi_i \cap f_{d-m}$ for $i \in J$. Then $\{\pi_i\}$ is the set of facets of $f = F_{d-m}$.

Let ϕ_1 be the lifting injection for F_{d-m-1} , in particular ϕ_1 takes $\{\Pi_i\}$ to $\Sigma_{r-1} \setminus A$, where A is the set of additional nodes. Denote by ψ the map $\pi_i \rightarrow \Pi_i$. Denote $\phi = \phi_1 \circ \psi$. Then ϕ is an injection from $\{\pi_i\}$ to $\Sigma \setminus A$. Let $\{\Pi_i \mid i \in I\}$ be a missing face of $D(F_{d-m})$ (where $I \subset J$ is some index set). By Prop. 1.8, either $\{\pi_i \mid i \in I\}$ or $\{f_{d-m}\} \cup \{\pi_i \mid i \in I\}$ is a missing face of $D(F_{d-m-1})$. This proves property 4) of the definition of liftings for either Σ_{r-1} or $\langle \Sigma_{r-1}, v_{d-m} \rangle$. By the same proposition, if $\{\Pi_i \mid i \in K\}$ is not a missing face in $D(F_{d-m})$ then neither $\{\pi_i \mid i \in K\}$ nor $\{f_{d-m}\} \cup \{\pi_i \mid i \in K\}$ is a missing face in $D(F_{d-m-1})$. This proves property 3).

Thus, either A or $A \cup \{f_{d-m}\}$ contains a set of additional nodes for some k -lifting of $D(F_{d-m-1})$. Hence, $k \leq d - (m + 1) + 1 = d - m$ and everything is proved. $\square$

Corollary 3.1. *Let $P \subset \mathbb{H}^d$ be a compact Coxeter polytope and f be a face of P . Then $D(f)$ contains no large missing faces. In particular, if $\dim f > 4$ then f is not a simplex.*

Proof. By the definition of liftings, for any missing face M of D there exists a Lannér diagram L in Σ such that $|L| \geq |M|$. Hence, the statement follows from the fact that a Lannér diagram contains at most 5 nodes. $\square$

Lemma 3.3. *Let D be a diagram of missing faces such that $|M| = 5$ for any $M \in \mathcal{M}_D$. Then any lifting of D is a 0-lifting.*

Proof. Suppose that D is a k -lifting, where $k > 0$, and ϕ is a lifting injection. Remove from D all additional nodes and denote by D_1 the diagram obtained. Then D_1 is a 0-lifting, and ϕ is a lifting bijection. Indeed, the conditions 2), 3) and 5) of the definition of k -lifting hold evidently.

For the condition 4), consider any missing face $M \in \mathcal{M}_D$. By definition, $\phi(M)$ belongs to some Lannér diagram of Σ . Note, that $\langle \phi(M) \rangle$ is a subdiagram of order 5. Since the order of a Lannér diagram does not exceed 5, $\langle \phi(M) \rangle$ is a Lannér diagram. $\square$

Now we will prove several lemmas about liftings we will use later.

Notation.

- Let D be a diagram of missing faces and $N_1, \dots, N_r \subset D$. We write $D = \bigcup_{i=1}^r N_i$ if for any node of D there exists a set $N_i, i \in \{1, \dots, r\}$ containing this node.

- Let $D = \bigcup_{i=1}^r N_i$ be a diagram of missing faces, where $N_i \cap N_j = \emptyset$ for any $i \neq j$. Suppose that there exists $k \in \{1, \dots, r\}$ such that $M \in \mathcal{M}_D$ if and only if $M = \bigcup_{t=i}^{i+k-1} N_t$ for some $i \in \{1, \dots, r-k+1\}$. Then we write $D = \lfloor N_1, N_2, \dots, N_r \rfloor_k$.
- When we are interested in combinatorial type of D only rather than in concrete subdiagrams N_i , we write $\lfloor |N_1|, \dots, |N_r| \rfloor_k$. For example, $\lfloor 1, 4, 1, 3 \rfloor_2$ stays for the diagram



- If there exists $k \in \{1, \dots, r\}$ such that $M \in \mathcal{M}_D$ if and only if $M = \bigcup_{t=i}^{i+k-1} N_t$ for some $i \in \{1, \dots, r\}$, where t is considered as a number modulo r , we write $D = (N_1, N_2, \dots, N_r)_k$ or $(|N_1|, \dots, |N_r|)_k$.
- We write $\Sigma \approx \lfloor N_1, N_2, \dots, N_r \rfloor_k$, if Σ contains no parabolic subdiagrams and the structure of Lannér subdiagrams of Σ corresponds to the diagram of missing faces $\lfloor N_1, N_2, \dots, N_r \rfloor_k$, i.e. Σ consists of disjoint subdiagrams $L_1, \dots, L_r$, $|L_i| = |N_i|$, and $L \subset \Sigma$ is a Lannér subdiagram if and only if $L = \langle L_i, L_{i+1}, \dots, L_{i+k-1} \rangle$ for some $i \in \{1, \dots, r-k+1\}$.
Similarly we use the notation $\Sigma \approx \lfloor |N_1|, \dots, |N_r| \rfloor_k$, $\Sigma \approx (N_1, N_2, \dots, N_r)_k$ and $\Sigma \approx (|N_1|, \dots, |N_r|)_k$.

Proposition 3.1 ([E1], Lemma 4.7). *Let $D = \lfloor M, N \rfloor_1$ be a diagram of missing faces, where $|M| \geq 3$ and $|N| \geq 4$. Then D has no 0-liftings.*

Lemma 3.4. *The diagram Θ_1 shown in Fig. 1 is the only lifting of $D = \lfloor 4, 5 \rfloor_1$. The additional node of this lifting is the leaf of Θ_1 which does not belong to a triple edge.*


Proof. Let Σ be a lifting of $D = \lfloor M_4, M_5 \rfloor_1$, where $|M_5| = 5$ and $|M_4| = 4$, and let ϕ be a lifting injection. Suppose that $\langle \phi(M_4) \rangle$ is a Lannér diagram of Σ . Then by Prop. 3.1 $\Sigma \approx \lfloor \phi(M_4), \phi(M_5) \rfloor_1$ is a superhyperbolic diagram. Hence, $\langle \phi(M_4) \rangle$ is an elliptic diagram and Σ contains at least one additional node a . Then the diagram $\langle a, \phi(M_4), \phi(M_5) \rangle$ satisfies the properties 1)-4) of the definition of liftings. So, by property 5) a is a unique additional node of Σ , and Σ is a 1-lifting. By property 3) of the definition of liftings, any Lannér subdiagram of Σ consists of 5 nodes. Since $|\Sigma| = |\langle a, \phi(M_4), \phi(M_5) \rangle| = 10$, Lemma 2.1 implies that Σ is one of the diagrams Θ_1 , Θ_2 and Θ_3 shown in Fig. 1. Note that Σ contains two disjoint Lannér diagrams $\langle \phi(M_4), a \rangle$ and $\langle \phi(M_5) \rangle$. Hence, $\Sigma = \Theta_1$ or Θ_2 .

It is easy to find a unique lifting injection for Θ_1 : a is the only leaf of Σ which does not belong to a triple edge, $\phi(M_4)$ are the remaining points in the lower row of vertices (see Fig. 1).

Lifting injection for Θ_2 does not exist: $\phi(M_4)$ belongs to any Lannér diagram different from $\phi(M_5)$, on the other hand, Θ_2 contains no such a quadruple of vertices, so the lemma is proved. $\square$

Lemma 3.5. *Let $D = N \cup \{x_1, x_2, x_3\}$ be a diagram of missing faces, where $|N| = 4$. If $\mathcal{M}_D = \{N \cup x_1, N \cup x_2, N \cup x_3\}$ then D has no liftings.*

Proof. Suppose that Σ is a lifting of D and ϕ is a lifting injection. By Lemma 3.3, Σ is a 0-lifting. Now, consider $\Sigma_{ij} = \langle \phi(N \cup \{x_i, x_j\}) \rangle$, $i \neq j$. Clearly, $\Sigma_{ij} \approx \lfloor 1, 4, 1 \rfloor_2$. By [E1, Lemma 5.3], $\lfloor 1, 4, 1 \rfloor_2$ is one of the following diagrams:



Taking each of these diagrams for Σ_{12} and trying to add x_3 in order to compose a correct diagram Σ_{13} , we obtain a conflicting subdiagram. $\square$

Lemma 3.6. *The diagram $D = [4, 4]_1$ has no 0-liftings, and any 1-lifting of D is one of the two diagrams shown in Fig.2.*




Figure 2: 1-liftings of $[4, 4]_1$.

Proof. By Prop. 3.1, there are no 0-liftings of D . Let Σ be a 1-lifting of D , let ϕ be a lifting injection, and let a be the additional node.

Denote by M_1 and M_2 the missing faces of D . We may assume that $\langle \phi(M_1) \rangle$ is an elliptic subdiagram of Σ and $\langle \phi(M_1), a \rangle$ is a Lannér diagram. We consider two cases:

Case 1. Suppose that $\langle \phi(M_2) \rangle$ is a Lannér diagram. By Prop. 3.1, Σ contains an additional Lannér diagram $L = \langle \phi(M_1), x \rangle$, where $x \in \phi(M_2)$. By Lemma 3.5, this additional Lannér diagram L is unique. Thus, $\Sigma \approx [a, \langle \phi(M_1) \rangle, x, \langle \phi(M_2) \setminus x \rangle]_2 = [1, 4, 1, 3]_2$. This is impossible by [E1, Folgerung 5.10].

Case 2. Suppose that both $\langle \phi(M_1) \rangle$ and $\langle \phi(M_2) \rangle$ are elliptic. Then $\langle \phi(M_1), a \rangle$ and $\langle \phi(M_2), a \rangle$ are Lannér diagrams. Note that a is an open vertex of $\langle \phi(M_1), a \rangle$, otherwise Σ has a conflicting Lannér diagram. Similarly, a is an open vertex of $\langle \phi(M_2), a \rangle$. Hence, each of $\langle \phi(M_1), a \rangle$ and $\langle \phi(M_2), a \rangle$ is one of $\mathcal{L}_1^5$ and $\mathcal{L}_5^5$.

If Σ contains an edge joining $\langle \phi(M_1) \rangle$ with $\langle \phi(M_2) \rangle$ then Σ has a conflicting subdiagram. Therefore, $\langle \phi(M_1) \rangle$ and $\langle \phi(M_2) \rangle$ are not connected in Σ , and Σ is one of the two diagrams shown in Fig. 2 (if $\langle \phi(M_1), a \rangle = \langle \phi(M_2), a \rangle = \mathcal{L}_5^5$ then Σ contains a parabolic subdiagram). $\square$

Denote by P_8 a unique compact hyperbolic Coxeter 8-polytope with 11 facets (see [Bu1],[E1]), and denote by $\Sigma(P_8)$ its Coxeter diagram, see Fig.3.




Figure 3: A unique compact hyperbolic Coxeter 8-polytope P_8 with 11 facets.

Lemma 3.7. *Let $D = \langle 4, 4, 2 \rangle_1$. Let Σ be a 0- or 1-lifting of D containing no Lannér diagram of order 3. Suppose that the positive inertia index of Σ does not exceed 8. Then $\Sigma = \Sigma(P_8)$.*

Proof. By Lemmas 3.1 and 3.6, D has no 0-liftings. Let Σ be a 1-lifting of D , let ϕ be a lifting injection, and let a be the additional node. Denote by M_1, M_2 and M_3 the missing faces of D , where $|M_1| = |M_2| = 4$ and $|M_3| = 2$. Let $M_3 = \{u_1, u_2\}$. By Lemma 3.6, $\langle \phi(M_1), \phi(M_2), a \rangle$ is one of the two diagrams shown in Fig.2. Since Σ has only one additional node, the absence of

Lannér diagram of order 3 implies that $\langle \phi(M_3) \rangle$ is a Lannér diagram. Therefore, any Lannér subdiagram of Σ distinct from $\langle \phi(M_3) \rangle$ is a Lannér diagram of order 5.

Suppose that the node u_1 does not belong to any additional Lannér subdiagram of Σ . Then $\langle \phi(M_1), \phi(M_2), u_1 \rangle$ is an elliptic subdiagram of Σ of order 9 in contradiction to the assumption that the positive inertia index of Σ is at most 8.


Hence, both u_1 and u_2 belong to some additional Lannér diagrams. Consider $X_1 = \langle u_1, \phi(M_1), \phi(M_2), a \rangle$. X_1 is a subdiagram of order 10 containing no small Lannér diagrams. Moreover, each vertex of X_1 belongs to some Lannér diagram. By Lemma 2.1, X_1 is one of the diagrams Θ_1 , Θ_2 and Θ_3 shown in Fig. 1. By the assumption of the lemma, the positive inertia index of Σ is at most 8. Therefore, $\det(X_1) = 0$ and $X_1 = \Theta_1$ (see Lemma 2.1). By Lemma 3.6, $\langle \phi(M_1), \phi(M_2), a \rangle$ is a linear subdiagram of Θ_1 . Hence, u_1 is the only leaf of Θ_1 that is not incident to a triple edge. Similarly, for $X_2 = \langle u_2, \phi(M_1), \phi(M_2), a \rangle$ we have $X_2 = \Theta_1$. Moreover, u_1 and u_2 are joined with different vertices of $\langle \phi(M_1), \phi(M_2), a \rangle$, otherwise one of the subdiagrams $\langle \phi(M_i), a, u_1, u_2 \rangle$ is superhyperbolic. Hence, $\Sigma = \Sigma(P_8)$. $\square$

Lemma 3.8. *Let $D = (1, 4, 1, 3, 1)_2$. Let Σ be a 0- or 1-lifting of D containing no Lannér diagram of order 3. Suppose that the positive inertia index of Σ does not exceed 8. Then $\Sigma = \Sigma(P_8)$.*

Proof. Let ϕ be a lifting injection. Denote the subsets of D as follows: $(1, 4, 1, 3, 1)_2 = (v, N_4, u, N_3, w)_2$.

Suppose $\langle \phi(N_3), \phi(u) \rangle$ is a Lannér diagram. Then $\langle \phi(v), \phi(N_4), \phi(u), \phi(N_3) \rangle \approx [1, 4, 1, 3]_2$, which is impossible by [E1, Folgerung 5.10]. Therefore, $\langle \phi(N_3), \phi(u) \rangle$ is an elliptic diagram and Σ is not a 0-lifting of D .

Now, assume that Σ is a 1-lifting and a is a unique additional node of Σ . Then the diagram $X = \langle a, \phi(D \setminus \{w\}) \rangle$ satisfies the conditions of Lemma 2.1. Since $|X| = 10$, X is one of the diagrams Θ_1 , Θ_2 and Θ_3 shown in Fig. 1. By the assumption, the positive inertia index of Σ is at most 8. Hence, $\det(X) = 0$ and $X = \Theta_1$:



Consider now $\langle X, \phi(w) \rangle$. Since $\langle \phi(N_4), \phi(w) \rangle$ should be an elliptic diagram, $\phi(w)$ does not attach to $\langle x_6, x_7, x_8, x_9 \rangle = \langle \phi(N_4) \rangle$. Further, $\phi(w)$ is not connected to $\langle x_1, x_2 \rangle$, otherwise $\langle x_1, x_2, x_3, \phi(w) \rangle$ is a conflicting subdiagram. So, besides $\phi(v)$, $\phi(w)$ may be attached to x_3 , x_4 , or x_5 only. Any edge joining w with x_3, x_4, x_5 is a simple edge, otherwise one of the diagrams $\langle x_2, x_3, x_4, \phi(w) \rangle$, $\langle x_3, x_4, x_5, \phi(w) \rangle$ and $\langle x_4, x_5, x_6, \phi(w) \rangle$ is a conflicting subdiagram. In order to obtain no parabolic subdiagrams, $\phi(w)$ should be connected to one of the vertices x_3, x_4, x_5 only (except $\phi(v)$).


If $\phi(w)$ attaches to x_3 then the subdiagram $\Sigma \setminus \phi(v)$ is superhyperbolic. If $\phi(w)$ is connected to x_5 then $\phi(N_3 \cup \{w\})$ is not contained in any Lannér diagram in contradiction to the definition of liftings. Thus, $\phi(w)$ attaches to x_4 . By assumption Σ contains no Lannér diagrams of order 3, so $\langle \phi(v), \phi(w) \rangle$ is a Lannér diagram and $\Sigma = \Sigma(P_8)$. $\square$

Lemma 3.9. *The diagram $D = [1, 3, 2, 3]_2$ has no 0-liftings.*

Proof. Suppose that Σ is a 0-lifting of $D = [1, 3, 2, 3]_2$. Comparing the 0-liftings of the diagram $[3, 2, 3]_2$ (see [E1, Lemma 5.12]) with the 0-liftings of the diagrams $[2, 3, 1]_2$ (see [T, Table 4, item 8]), we obtain that Σ is one of the following diagrams:

$$\Sigma \approx [\langle \phi(M_5) \setminus \{x_1, x_2\} \rangle, \langle x_1, x_2 \rangle, \langle \phi(M_3) \rangle, a]_2 = [3, 2, 3, 1]_2,$$

which is impossible by Lemma 3.9. Hence, Σ contains at least one more additional Lannér diagram L' . Suppose that $L' = \langle \phi(M_3), x_1, x_3 \rangle$, $x_3 \in \phi(M_5)$, $x_3 \neq x_2$. As it was shown above, the open vertex x_1 of $\phi(M_5)$ is connected with two other nodes, x_2 and x_3 , so $\langle \phi(M_5) \rangle = \mathcal{L}_5^5$. In particular, the edges x_1x_2 and x_1x_3 are simple. Now, taking in mind that $\langle L, a \rangle \approx [a, \langle \phi(M_3) \rangle, \langle x_1, x_2 \rangle]_2 = [1, 3, 2]_2$ and $\langle L', a \rangle \approx [a, \langle \phi(M_3) \rangle, \langle x_1, x_3 \rangle]_2 = [1, 3, 2]_2$, we check all possibilities for $[1, 3, 2]_2$ such that x_1x_2 and x_1x_3 are simple edges (see [T, Table 4, item 8]). In this way we obtain that either Σ contains some conflicting Lannér diagram or Σ is one of the following diagrams:



All these diagrams are superhyperbolic. We come to a contradiction which proves the lemma. $\square$

4 Absence of polytopes in dimensions ≥ 9

Suppose that there exists a compact hyperbolic Coxeter d -polytope P with $d+4$ facets. Denote by Σ its Coxeter diagram.

We will refer to the classification of compact hyperbolic Coxeter d -polytopes with $d+k$ facets for $k \leq 3$ (see [L], [E2], [K] and [T]). In particular, we remind that P_8 is a unique Coxeter 8-polytope with 11 facets, and $\Sigma(P_8)$ is its Coxeter diagram (see Fig. 3).

From now on by polytope we mean a compact hyperbolic Coxeter polytope. By “a diagram of a polytope” we mean a Coxeter diagram of the polytope.

Lemma 4.1. *Suppose that $d \geq 9$. Then*

- (1) *any node of Σ is incident to at most one non-simple edge;*
- (2) *Σ contains no Lannér diagram of order 3;*
- (3) *Σ contains no edge of multiplicity ≥ 4 .*

Proof. Suppose the lemma is broken. Then there exists either a node incident to two dotted edges, or a subdiagram $S_0 \subset \Sigma$ of the type G_2^m for some $m \geq 4$ having at least one bad neighbor.

Suppose that there exists a node v incident to two dotted edges. Then v corresponds to a $(d-1)$ -dimensional (non-Coxeter) face f with at most $d+1 = (d-1) + 2$ facets, so f is either a simplex or a product of two simplices. If f is a simplex then P has a large missing face, which is impossible by Cor. 3.1. If f is a product of two simplices and it has no large missing faces then f is a product of two 4-dimensional simplices and $D(f) = [5, 5]_1$ (since $d \geq 9$). By Prop. 3.1 the diagram $[5, 5]_1$ has no liftings, so we contradict Lemma 3.2.

Suppose that Σ contains a subdiagram $S_0 \subset \Sigma$ of the type G_2^m for some $m \geq 4$ having at least one bad neighbor. Then $P(S_0)$ is a Coxeter $(d-2)$ -polytope with at most $(d-2) + 3$ facets, so $(d-2)$ is either equal to 8 or less than 7 (see [T]). If $d-2 < 7$ we obtain that $d \leq 8$. If $d-2 = 8$ we have $d = 10$, and $P(S_0)$ is a unique Coxeter 8-polytope with 11 facets (see Fig. 3). The diagram Σ_{S_0} contains a subdiagram of the type H_4 with 2 neighbors (the neighbors are attached to this diagram by simple edges). Cor. 1.2 implies that this diagram corresponds to a subdiagram $S_1 \subset \Sigma$ of the type H_4 with 2 neighbors. So, $P(S_1)$ is a Coxeter 6-polytope with at most 8 facets, which is impossible (see [E2], [K]). $\square$

Lemma 4.2. *Suppose that $d \geq 9$. Then Σ does not contain any Lannér diagram of order 5.*

Proof. Suppose there exists a diagram $L_0 \subset \Sigma$ of order 5. Clearly, it contains a subdiagram $S_0 \subset L_0$, such that $S_0 = H_4$ or F_4 . Then $P(S_0)$ is a $(d-4)$ -dimensional Coxeter polytope, and its Coxeter diagram Σ_{S_0} is a subdiagram of Σ (Cor. 1.1). The diagram S_0 has at least one neighbor, the node $L_0 \setminus S_0$. Thus, $P(S_0)$ has either $d-3$, or $d-2$, or $d-1$ facets.

Case 1: $P(S_0)$ has $d-3$ facets. Then $P(S_0)$ is a Coxeter simplex, and we obtain $d-4 \leq 4$, i.e. $d \leq 8$.

Case 2: $P(S_0)$ has $d-2$ facets. Then $P(S_0)$ is a Coxeter $(d-4)$ -polytope with $(d-4)+2$ facets. Results of [E1] and [K] imply that $d-4 \leq 5$, i.e. $d \leq 9$. If $d = 9$, $P(S_0)$ is a 5-dimensional prism. Since $\Sigma_{S_0} = \overline{S}_0$, condition (1) of Lemma 4.1 and [K] imply that there are only four possibilities for Σ_{S_0} : all of them are presented on Fig. 4.





Figure 4: 5-prisms satisfying condition (1) of Lemma 4.1.

Let L_1 be a unique Lannér subdiagram of $\overline{S}_0$ of order 5, and denote by S_1 a unique subdiagram of L_1 of the type H_4 or F_4 . Denote $u = L_1 \setminus S_1$. The diagram consists of the following parts:




where a_1 and a_2 attach to S_0 , and S_0 is not connected to $\overline{S}_0$. Since any two indefinite subdiagrams of Σ are joined in Σ by at least one edge, each of a_1 and a_2 attaches to L_1 . If they both attach to S_1 , then S_1 has at least 3 neighbors and we refer to Case 1. So, one of a_1 and a_2 , say a_1 , attaches to $u = L_1 \setminus S_1$ (see Fig. 4) and does not attach to S_1 .


Consider the diagrams (a) and (b) with nodes indexed as shown in Fig. 4. Let $S_2 = \overline{S}_0 \setminus \langle u_2, u_4 \rangle$. S_2 is a diagram of type B_5 with at least 3 bad neighbors (u_4 , u_2 and a_1). Thus, a_2 does not attach to S_2 . Considering a subdiagram $S_3 = \overline{S}_0 \setminus \langle u_2, u_3 \rangle$ instead of S_2 , we obtain that a_2 does not attach to u_4 either. Thus, a_2 does not attach to L_1 , and an indefinite subdiagram $\langle S_0, a_2 \rangle$ is not joined with L_1 . This means that Σ is superhyperbolic, which is impossible.

Now let $\bar{S}_0$ be one of the diagrams (c) and (d) with nodes indexed as shown in Fig. 4. We have two possibilities: a_2 either attaches to S_1 or not. Suppose that a_2 does not attach to S_1 . Then $P(S_1)$ is a Coxeter 5-polytope with 8 facets. However, the Coxeter diagrams of such polytopes do not satisfy Lemma 4.1 (see [T]). Thus, a_2 is joined with S_1 and the diagram S_1 has exactly two neighbors (otherwise we refer to Case 1), so the polytope $P(S_1)$ is a Coxeter 5-prism. By the reasoning of the previous paragraph we may assume that the Coxeter diagram Σ_{S_1} of $P(S_1)$ is of the type (c) or (d) (see Fig. 4). Notice that the diagram $\bar{S}_1 = \Sigma_{S_1}$ contains S_0 . Therefore, S_0 is of the type H_4 .

Further, the diagram $\bar{S}_1$ contains u_1 , u_2 and a_1 . Since $P(S_1)$ is a 5-prism and $\bar{S}_1 = \langle S_0, u_1, u_2, a_1 \rangle$, $\bar{S}_1$ contains a Lannér diagram of order 5. The nodes u_1 and u_2 do not attach to S_0 , so $\langle S_0, a_1 \rangle$ is a Lannér diagram. It follows that it should be joined with L_1 , and by Lemma 1.4 a_1 attaches to u by a dotted edge. Therefore, the diagram $\langle S_0, a_1, u, S_1 \rangle$ is



Recall that the nodes u_1 and u_2 are joined by a dotted edge. Since $u_1, u_2 \in \bar{S}_0$ and $u_1, u_2 \in \bar{S}_1$, they may be joined in $\Sigma \setminus a_2$ with u and a_1 only. We obtain the following four possibilities for the diagram $\Sigma \setminus a_2$:



By the assumption, the node a_2 attaches to $S_1 = \langle v_5, v_6, v_7, v_8 \rangle$. Since $a_2 \notin \Sigma_{S_0}$, a_2 also attaches to $S_0 = \langle v_1, v_2, v_3, v_4 \rangle$. Further, a_2 attaches to $\langle u_1, u_2 \rangle$, otherwise an indefinite diagram $\langle S_0, a_2 \rangle$ is not joined with the Lannér diagram $\langle u_1, u_2 \rangle$. Since the diagrams $\langle v_2, v_3, v_4, a_1, u_1 \rangle$, $\langle v_7, v_6, v_5, u, u_1 \rangle$ and $\langle v_7, v_6, v_5, u, u_2 \rangle$ already have 3 bad neighbors each, no dotted edge ends in a_2 (see (1) of Lemma 4.1). Carefully examining possible multiplicities of edges, it is easy to see that we always obtain either a Lannér subdiagram of order 3, or a parabolic subdiagram, or a subdiagram of the type H_4 with at least 3 neighbors, and we refer to Case 1.

Case 3: $P(S_0)$ has $d - 1$ facets. Then $P(S_0)$ is a Coxeter $(d - 4)$ -polytope with $(d - 4) + 3$ facets. Since $d \geq 9$, we have $d - 4 \geq 5$. According to [T], Coxeter diagrams of all such polytopes either do not satisfy the conditions of Lemma 4.1, or contain a subdiagram of the type H_4 with at least two neighbors, so we refer to the previous cases. $\square$


Lemma 4.3. *Suppose that one of the following holds:*

- 1) $d \geq 9$, or
 - 2) $d = 8$, Σ satisfies conditions (1)–(3) of Lemma 4.1 and Σ does not contain any Lannér diagram of order 5.
- Then Σ does not contain any Lannér diagram of order 4.*

Proof. Suppose the contrary. Let L_0 be a Lannér subdiagram of Σ of order 4. Denote by S_0 a subdiagram of L_0 of the type H_3 (if any) or B_3 (otherwise). Then $P(S_0)$ is a Coxeter $(d - 3)$ -polytope with at most $(d - 3) + 3$ facets, and its Coxeter diagram Σ_{S_0} does not contain

Lannér diagrams of order 5 (Lemmas 4.2 and 3.2). If $P(S_0)$ is a simplex or a product of two simplices, then $d - 3 \leq 3$ or $d - 3 \leq 4$ respectively, so in this case $d \leq 7$.

Suppose that $P(S_0)$ has $(d - 3) + 3$ facets. According to [T], for $d \geq 8$ the Coxeter diagrams of almost all Coxeter $(d - 3)$ -polytopes with d facets contain Lannér diagrams of order 5. There is only one exclusion: a 5-polytope with 8 facets having the following Coxeter diagram Σ_{S_0} :



By Cor. 1.2, no node of $\overline{S_0} \setminus \langle b_1, b_2 \rangle$ is a good neighbor of S_0 . The node b_1 is a good neighbor of S_0 , otherwise Σ contains an order 3 Lannér diagram $\langle b_1, c_1, d_1 \rangle$, which contradicts the assumption. Thus, the edge b_1c_1 is a simple edge of Σ , and the subdiagram $\langle d_1, c_1, b_1, x_1, x_2 \rangle \subset \Sigma$ is a Lannér diagram of order 5 (where x_1 and x_2 are the ends of the simple edge of S_0). The contradiction with the assumption of the lemma completes the proof. $\square$

Theorem 1. *There is no compact hyperbolic Coxeter d -polytope with $d + 4$ facets for $d \geq 9$.*

Proof. Lemmas 4.1, 4.2 and 4.3 imply that all Lannér diagrams in Σ have order 2. By [E1, Satz 6.9], in this case $d \leq 4$. $\square$

5 Absence of polytopes in dimension 8

In this section, we prove that no compact Coxeter polytope with 12 facets exists in $\mathbb{H}^8$. Suppose there exists an 8-polytope P with 12 facets. First, we show some properties of its Coxeter diagram Σ . Most of them repeat ones that hold for polytopes in large dimensions. However, the proofs in eight-dimensional case are much more complicated.

Lemma 5.1 ([FT2], Lemma 1). *Let $\Sigma(P)$ be a Coxeter diagram of a Coxeter d -polytope P . Then no proper subdiagram of $\Sigma(P)$ is a Coxeter diagram of a finite volume Coxeter d -polytope.*

In particular, Σ does not contain $\Sigma(P_8)$ as a proper subdiagram.

Lemma 5.2. *Suppose that nodes v_1 and v_2 of Σ are joined by an edge of multiplicity $k \geq 2$. Then*

- (1) *the subdiagram $\langle v_1, v_2 \rangle$ has at most one bad neighbor;*
- (2) *$k \leq 3$;*
- (3) *if the subdiagram $\langle v_1, v_2 \rangle$ has a bad neighbor, then it has at least one good neighbor, too.*

Proof. Denote by S_0 the diagram $\langle v_1, v_2 \rangle$. If S_0 has at least two bad neighbors, $P(S_0)$ is a Coxeter 6-polytope with at most 8 facets, which is impossible [K], [E1].

Suppose $k \geq 4$ or S_0 has a unique neighbor which is bad. In particular, all neighbors of S_0 are bad. In this case $P(S_0)$ is a Coxeter 6-polytope with at most 9 facets, and its Coxeter diagram Σ_{S_0} is a subdiagram of Σ . By [T], there are only three such polytopes, their diagrams are shown in Fig. 5. A Coxeter diagram of each of these three polytopes contains a multiple edge with at least two bad neighbors (see Fig. 5). $\square$




Figure 5: Compact hyperbolic Coxeter 6-polytopes with 9 facets.

Lemma 5.3. (1) Σ contains no Lannér diagram of order 3.
(2) No end of a dotted edge is incident to a triple edge.
(3) If Σ contains a unique dotted edge, then no end of that edge is incident to a double edge.

Proof. Suppose that the lemma is broken. Denote by L_0 a Lannér diagram of order 3 or a subdiagram consisting of adjoint dotted and multiple edges. L_0 contains an edge of multiplicity 2 or 3. We consider the following two cases:

Case 1: L_0 contains an edge of multiplicity 3.

Denote by S_0 the subdiagram consisting of this edge. $P(S_0)$ is a Coxeter 6-polytope with at most 9 facets, so we may assume that $P(S_0)$ has exactly 9 facets. Then Σ_{S_0} is one of the diagrams shown in Fig. 5. By Cor. 1.1, the diagram $\overline{S}_0 \subset \Sigma$ can be obtained from Σ_{S_0} by replacing some dotted edges by ordinary edges, and some edges labeled by 10 by simple edges. Applying (or not) such a procedure to the Coxeter diagrams of the polytopes P_6^1 and P_6^3 we see that the resulting diagrams contain either a parabolic subdiagram or a multiple edge with at least two bad neighbors.

Now suppose that $P(S_0) = P_6^2$. Denote by u a unique bad neighbor of S_0 (it is unique by Lemma 5.2). By Cor. 1.1, Σ_{S_0} can be transformed to $\overline{S}_0$ by the change of some edges labeled by 10 into simple edges and some dotted edges into ordinary edges. By Cor. 1.2, the dotted edge of Σ_{S_0} remains dotted in $\overline{S}_0$. By Lemma 5.2, the edge of Σ_{S_0} labeled by 10 is a simple edge of $\overline{S}_0$. Therefore, the leaf $a \in \Sigma_{S_0}$ that is incident to the edge labeled by 10 is a good neighbor of S_0 in Σ . The remaining nodes of Σ_{S_0} cannot be good neighbors of S_0 in view of Cor. 1.2. So, the subdiagram $\Sigma \setminus u$ is a Coxeter diagram $\Sigma(P_8)$, which is impossible by Lemma 5.1.

Case 2: L_0 contains no edge of multiplicity 3.

Denote by S_0 the subdiagram consisting of a double edge contained in L_0 . $P(S_0)$ is again a 6-polytope with 9 facets. By Prop. 1.11, the diagram $\overline{S}_0 \subset \Sigma$ can be obtained from Σ_{S_0} by replacing some dotted edges by ordinary (or empty) edges, and some double edges by simple or empty edges. Applying (or not) such a procedure to the Coxeter diagrams of the polytopes P_6^3 and P_6^2 we see that the resulting diagrams contain a multiple edge (a triple one or labeled by 10 respectively) with at least two bad neighbors.

Suppose that $P(S_0) = P_6^1$. Both double edges from Σ_{S_0} become simple or empty edges in $\overline{S}_0$, otherwise Σ contains a Lannér diagram with a triple edge and we refer to Case 1. Furthermore, by Cor. 1.2, only one end of each double edge can be a good neighbor of S_0 . Hence, by Prop. 1.11 both double edges of Σ_{S_0} are simple edges in $\overline{S}_0$. This implies that a unique dotted edge in Σ_{S_0} remains a dotted edge in $\overline{S}_0$ (otherwise the diagram $\overline{S}_0$ is superhyperbolic). So, in this case we may assume that L_0 contains no dotted edge. By Prop. 1.11, there exist only two good neighbors of S_0 — the ends of the dotted edge. They attach to S_0 by simple edges. Consider

the diagram S_1 of the type H_4 containing a triple edge, a simple edge that was double in Σ_{S_0} and a simple edge joining S_0 with its good neighbor. Since L_0 is a Lannér diagram without edges of multiplicity greater than two, L_0 contains 3 edges, and we obtain that the diagram S_1 of the type H_4 has at least 4 bad neighbors (two neighbors in $\overline{S}_0$ and two ones in S_0). This contradicts lemma 1.3, which completes the proof of the lemma. $\square$

The following lemma is a corollary of the main result of [FT2].

Lemma 5.4 ([FT2], cor. of Th. A). *If $\Sigma(P)$ is a Coxeter diagram of a compact Coxeter polytope $P \subset \mathbb{H}^d$, where $d > 4$, then $\Sigma(P)$ contains a dotted edge.*

Lemma 5.5. *Let $S \subset \Sigma$ be an elliptic subdiagram of order 3 containing no component of the type A_n . Then S has at most 2 bad neighbors.*

Proof. Suppose that S has 3 or more bad neighbors. Then Σ_S is a Coxeter 5-polytope with at most 6 facets, which is impossible. $\square$


Lemma 5.6. *Suppose that Σ contains a Lannér subdiagram L_0 of order 5 and that the dotted edge is unique in Σ . Then Σ contains a subdiagram S of the type F_4 or H_4 such that $P(S)$ is not a simplex.*

Proof. Suppose that the lemma is broken. Then we may assume that for any diagram $S' \subset \Sigma$ of the type H_4 or F_4 the face $P(S')$ is a Coxeter simplex.

Denote by S_0 a subdiagram of L_0 of the type H_4 or F_4 . In particular, $P(S_0)$ is a Coxeter 4-simplex. The Coxeter diagram Σ_{S_0} of $P(S_0)$ is a Lannér diagram of order 5. Denote it by L_1 . Denote by S_1 a subdiagram of L_1 of the type H_4 or F_4 . $P(S_1)$ is a Coxeter simplex, and its Coxeter diagram $\overline{S}_1 = \Sigma_{S_1}$ is a Lannér diagram of order 5. Denote it by L_2 . Notice that $S_0 \subset L_2$, so let $S_2 = S_0$. Let $v_i = L_i \setminus S_i$, $i = 0, 1, 2$.

Since $S_2 = S_0$ does not attach to L_1 , v_2 is joined with L_1 . On the other hand, L_2 does not attach to S_1 . Thus, the subdiagram $\langle L_1, L_2 \rangle$ consists of two Lannér diagrams L_1 and L_2 joined by a unique edge $v_1 v_2$, such that $L_i \setminus v_i$ are of the type H_4 or F_4 . By Lemma 1.4, $v_1 v_2$ is a dotted edge.

Now suppose the dotted edge $v_1 v_2$ is unique. Denote by a and b two nodes not contained in L_1 and L_2 . Note that if $v_2 \neq v_0$ then either a or b coincides with v_0 . Both a and b are neighbors of S_1 and S_2 . The diagram Σ consists of the following parts:



where a wave line means that the vertex is joined with the subdiagram by some non-dotted edges, and some additional non-dotted edges may appear in the subdiagram $\langle a, b, v_1, v_2 \rangle$.

By Lemma 5.3, vertices v_1 and v_2 attach to S_1 and S_2 respectively by simple edges only. Thus, L_1 (as well as L_2) is one of the diagrams $\mathcal{L}_1^5$, $\mathcal{L}_4^5$ or $\mathcal{L}_5^5$ (see Table 2).

Suppose that $L_1 = \mathcal{L}_5^5$. Let $w_1 \in S_1$ be a vertex attached to a . If $w_1 a$ is a triple edge, we obtain either a Lannér diagram of order 3 or a diagram of the type H_3 with at least three bad neighbors, which is impossible by Cor. 3.1. If $w_1 a$ is a double edge or a simple edge, we obtain

either a parabolic diagram or a Lannér diagram of order 3. So, the multiplicity of the edge w_1a should be greater than 3, which contradicts Lemma 5.2.

Thus, both S_1 and S_2 are of the type H_4 . For each of 4 possible pairs of diagrams L_1 and L_2 there exists a unique label of the dotted edge v_1v_2 such that the determinant of the diagram $\Sigma \setminus \langle a, b \rangle$ vanishes. The label equals $(1 + \sqrt{5})/2$ for $L_1 = L_2 = \mathcal{L}_1^5$, $4 + 2\sqrt{5}$ for $L_1 = L_2 = \mathcal{L}_4^5$, and $(3 + \sqrt{5})/\sqrt{2}$ for $L_1 = \mathcal{L}_1^5$, $L_2 = \mathcal{L}_4^5$ (or $L_2 = \mathcal{L}_1^5$, $L_1 = \mathcal{L}_4^5$).

Denote by $u_1 \in S_1$ and $u_2 \in S_2$ the leaves of $\Sigma \setminus \langle a, b \rangle$ incident to the triple edges. If we assume that each of a and b attaches to $S_1 \setminus u_1$ and to $S_2 \setminus u_2$, all edges joining them with $S_i \setminus u_i$ are simple, and a is not joined with b , then the diagram $\Sigma \setminus \langle v_1, v_2 \rangle$ contains a parabolic subdiagram of the type $\tilde{A}_m$ for some m , $2 \leq m \leq 7$. If in addition there are some multiple edges joining a and b with $S_1 \setminus u_1$ and to $S_2 \setminus u_2$, or a is joined with b , then $\Sigma \setminus v_1$ or $\Sigma \setminus v_2$ contains a subdiagram of the type H_3 with at least 3 bad neighbors, or a Lannér diagram of order 3, or a parabolic diagram of the type $\tilde{B}_3$. Therefore, at least one of a and b , say a , can attach to one of $S_i \setminus u_i$, say $S_1 \setminus u_1$, by multiple edges only. This means that we have got two cases: a either attaches to $S_1 \setminus u_1$ by a unique multiple edge (Lemma 5.3) or does not attach to $S_1 \setminus u_1$ (in the latter case a attaches to u_1).

Case 1: a attaches to $S_1 \setminus u_1$ by a unique multiple edge au . It is easy to see that in this case $L_1 = \mathcal{L}_4^5$, u is the leaf of L_1 different from v_1 , and a does not attach to v_1 (otherwise we obtain either a diagram of the type H_3 with at least 3 bad neighbors, or a Lannér diagram of order 3, or a parabolic subdiagram). If au is a double edge then $\langle a, L_1 \setminus u_1 \rangle$ is a parabolic diagram $\tilde{B}_4$. Suppose that au is a triple edge. The vertex a is not joined with u_1 , otherwise the diagram $\langle u, a, u_1 \rangle$ of the type H_3 has at least 3 bad neighbors, which contradicts Cor. 3.1. Now consider the diagram $\Sigma_1 = \langle L_1, a, v_2 \rangle$. The vertex a may attach (by simple edge only, see Lemma 5.3) to v_2 only. An elementary computation shows that Σ_1 is superhyperbolic whenever a attaches to v_2 or not.

Case 2: a does not attach to $L_1 \setminus u_1$. Therefore, a attaches to u_1 by a simple edge. Again, consider the diagram $\Sigma_1 = \langle L_1, a, v_2 \rangle$. The vertex a may attach (by simple edge only) to v_1 and v_2 only. For each of 4 pairs of diagrams L_1 and L_2 the diagram Σ_1 is superhyperbolic whenever a attaches to v_1 and (or) v_2 or not. □

Lemma 5.7. *Any node of Σ is incident to at most one dotted edge.*

Proof. Suppose the contrary. Denote by u a node of Σ incident to at least two dotted edges. Since the number of bad neighbors of u cannot exceed 3, u is incident to either two or three dotted edges. If there are three, the polytope $P(u)$ is a simplicial 7-face of P , which contradicts Cor. 3.1 of Lemma 3.2.

Thus, u is incident to exactly two dotted edges, and $P(u)$ is a 7-polytope with 9 facets, i.e. a polytope whose diagram of the missing faces is $[k, 9 - k]_1$ for some $k < 9$. Since the missing faces diagram of such a polytope contains no large missing faces, it is of the type $[4, 5]_1$. By Lemma 3.4, this diagram of missing faces has a unique lifting, namely the diagram Θ_1 shown in Fig. 1. Denote the nodes of Θ_1 as shown in Fig. 6, denote by v and w the remaining two nodes of Σ . Each of them attaches to the node u by a dotted edge and attaches to the subdiagram $\langle v_5, \dots, v_9 \rangle \subset \Sigma$, where the nodes are indexed as shown in Fig. 6.

Case 1: no dotted edge joins $\langle v, w \rangle$ with $\Sigma \setminus \langle v, w, u \rangle = \langle v_1, \dots, v_9 \rangle$.

Let $S_0 = \langle v_2, \dots, v_8 \rangle$. The face $P(S_0)$ is one-dimensional, so S_0 is contained in exactly two elliptic diagrams of order 8. This means that one of v and w , say v , either is not joined with S_0 or is a good neighbor of S_0 . In particular, any double edge from v to S_0 may end in v_2 or v_8 only. Now Lemma 5.3 implies that no double edge joins v with $\Sigma \setminus \langle v, w, u \rangle$. Thus, v may be joined by simple edges only with v_1, v_9 and one node of S_0 . A straightforward calculation




Figure 6: A unique lifting of $[4, 5]_1$.

shows that none of obtained subdiagrams $\Sigma \setminus \langle w, u \rangle$ is admissible with positive inertia index at most 8.

Case 2: at least one of v and w , say v , is joined by a dotted edge with some node $v_x \in \Sigma \setminus \langle v, w, u \rangle$.

In this case $P(v)$ is a 7-face of P with 9 facets, i.e. a face with a diagram of missing faces of the type $[5, 4]_1$. A unique lifting of its missing faces is the diagram Θ_1 (Lemma 3.4). In other words, diagram $\Sigma \setminus \langle u, v_x \rangle$ looks as shown in Fig. 6, where v takes place of u , w takes place of v_x , and, possibly, the node v is not joined with v_4 , but joined with either v_6 (if $v_x \neq v_6$) or with w (if $v_x = v_6$).

If v_x and w are not joined by a dotted edge, we obtain a parabolic diagram $\langle v_{x-1}, v_x, v_{x+1}, w \rangle$ (if $v_x \neq v_1, v_9$) or a Lannér diagram of order 3 (otherwise). Thus, v_x and w are joined by a dotted edge. Now consider $P(v_x)$. It is again a 7-polytope with 9 facets, but in the lifting of its missing faces diagram the node v_x is joined with at least two other nodes, which contradicts Lemma 3.2. □


Lemma 5.8. Σ contains at least two dotted edges.

Proof. By Lemma 5.4, there exists at least one dotted edge. Suppose it is unique. Then, by Lemmas 5.2 and 5.3, Σ satisfies conditions (1)–(3) of Lemma 4.1. Below we show that this implies that no Lannér diagram of order 5 is contained in Σ . In view of Lemmas 4.3, 5.3 and 1.2, this proves the lemma.

We assume that there exists a Lannér diagram $L_0 \subset \Sigma$ of order 5. Denote by S_0 a subdiagram of L_0 of the type H_4 or F_4 . The Coxeter diagrams of all Coxeter 4-polytopes with 7 facets contain either a Lannér diagram of order 3 or at least two dotted edges. The Coxeter diagrams of all Coxeter 4-polytopes with 6 facets that are not prisms contain Lannér diagrams of order 3. Thus, $P(S_0)$ is either a Coxeter 4-prism or a Coxeter 4-simplex.


By Lemma 5.6, we can assume that $P(S_0)$ is a prism. Consider two cases.

Case 1: $P(S_0)$ is a prism whose Coxeter diagram Σ_{S_0} is different from



where the label “2, 3” means that the nodes are either not joined or joined by a simple edge.


Looking at the list of Coxeter diagrams of 4-prisms [K], we see that $\Sigma_{S_0} = \overline{S}_0$ has a subdiagram S_1 of the type H_4 or F_4 with a dotted edge attached to S_1 . Since Σ contains a unique dotted edge, the Coxeter diagram $\Sigma_{S_1} = \overline{S}_1$ of $P(S_1)$ does not contain dotted edges, so $P(S_1)$ is a simplex. Notice that $S_0 \subset \overline{S}_1$. Denote by u and v the neighbors of S_0 , denote by a the end of the dotted edge not contained in S_1 , and by b the remaining node of Σ not contained in $\langle S_0, S_1 \rangle$. So, the diagram Σ consists of the following parts:




where u and v attach to S_0 , and S_0 does not attach to $\overline{S}_0$. Since $P(S_1)$ is a simplex, b and one of u and v , say v , attach to S_1 , and $\langle S_0, u \rangle = \overline{S}_1$ is a Lannér diagram.

Consider the diagram $\Sigma \setminus \langle v, a \rangle$. It consists of a Lannér diagram $\langle S_0, u \rangle = \overline{S}_1$ and an indefinite diagram $\langle S_1, b \rangle$. These two subdiagrams may be joined by a unique non-dotted edge ub only. It is easy to see that the diagram $\Sigma \setminus \langle v, a \rangle$ is superhyperbolic.

Case 2. $P(S_0)$ is a prism whose Coxeter diagram Σ_{S_0} is



Denote by S_1 a subdiagram of Σ_{S_0} of the type B_4 , say $S_1 = \langle v_1, v_2, v_3, v_5 \rangle$. So, Σ consists of the following parts:



where u and v attach to S_0 , and S_0 does not attach to $\overline{S}_0$. Notice that S_1 has at least two bad neighbors, v_4 and v_6 , so $P(S_1)$ is either an Esselmann polytope, or a prism, or a simplex. Consider these three cases.

Case 2.1: $P(S_1)$ is an Esselmann polytope.

Then Σ_{S_1} contains two disjoint Lannér diagrams of order 3, while $\overline{S}_1$ contains no subdiagram of order 3. Therefore, Σ_{S_1} contains two nodes that are good neighbors of the diagram S_1 in Σ . This is impossible by Cor. 1.2 (we use the list of Esselmann diagrams [E2] and the fact that S_1 is a diagram of the type B_4).

Case 2.1: $P(S_1)$ is a prism.

As above, u and v are neighbors of S_0 . By the assumption, Σ contains a unique dotted edge, hence at least one of u and v , say v , should be a good neighbor of S_1 . Then $\langle S_1, v \rangle$ is of the type B_5 . If u is also a good neighbor of S_1 , we obtain either a parabolic subdiagram of Σ or a Lannér diagram of order 3. Thus, v is a unique good neighbor of S_1 .

Consider the diagram $\overline{S}_1 = \langle S_0, u, v \rangle$. The only node of $\overline{S}_1$ that attaches to S_1 is v . Thus, $\overline{S}_1$ may differ from Σ_{S_1} by multiplicities of edges incident to v only. By Prop. 1.11, any simple edge of $\overline{S}_1$ incident to v becomes a double edge in Σ_{S_1} , and any other edge of $\overline{S}_1$ incident to v becomes a dotted edge. Since $P(S_1)$ is a prism, Σ_{S_1} contains a unique dotted edge. At the same time, no Coxeter diagram of a Coxeter 4-prism contains a node incident to a dotted edge and to a multiple edge simultaneously. Thus, v is joined with a unique node of $\overline{S}_1 \setminus v = \langle S_0, u \rangle$, say w , and $\overline{S}_1$ may be obtained from Σ_{S_1} by replacing the dotted edge by a double or a triple edge.

Recall that v is a unique neighbor of S_1 contained in $\overline{S}_1$, and it is joined with v_5 by a simple edge. Thus, we obtain either a parabolic subdiagram $\langle v_1, v_2, v_3, v_5, v, w \rangle$ or a subdiagram $\langle v_3, v_5, v, w \rangle$ of the type H_4 with at least 4 neighbors.

Case 2.3: $P(S_1)$ is a simplex.


We preserve the notation from Case 1. The difference is that now one of u and v , say v , is a bad neighbor of S_1 . The only way to attach a bad neighbor to S_1 such that no parabolic or Lannér diagram of order 3 appears is to join v by a triple edge with v_5 . But in this case we obtain again a subdiagram $\langle v_2, v_3, v_5, v \rangle$ of the type H_4 with at least 4 neighbors.

Hence, no Lannér subdiagram of Σ of order five exists, and the lemma is proved. $\square$

Theorem 2. *There are no compact hyperbolic Coxeter 8-polytopes with 12 facets.*

Proof. Suppose there exists a polytope P with a Coxeter diagram Σ . By Lemma 5.4, Σ contains a dotted edge vw . Lemma 5.7 implies that $P(v)$ is a 7-polytope with 10 facets. By Lemma 5.8, $P(v)$ has a pair of disjoint facets.

There are only four two-dimensional Gale diagrams of order 10 containing a missing face of order 2 and not containing large missing faces, all of them are listed below.



By Lemma 3.2, Σ contains either 0- or 1-lifting of one of these diagrams. By Lemma 5.3, Σ contains no Lannér diagram of order 3. Since P is an 8-dimensional polytope, the positive inertia index of Σ is at most 8. Hence, by Lemmas 3.7, 3.8, 3.10 and 3.11, Σ contains $\Sigma(P_8)$, which contradicts Lemma 5.1. $\square$

6 Polytopes in dimension 7

In this section we assume that Σ is the Coxeter diagram of a compact Coxeter 7-polytope with 11 facets and prove that Σ coincides with Σ_{P_7} , where Σ_{P_7} is a diagram found by V. Bugaenko in [Bu1] and shown in Fig. 7




Figure 7: A unique compact Coxeter 7-polytope with 11 facets.

Theorem 3. *If Σ is the Coxeter diagram of a compact Coxeter 7-polytope with 11 facets then $\Sigma = \Sigma_{P_7}$.*

The proof is similar to the proof in larger dimensions, however it is rather long and routine, with many cases to consider. First, we prove that Σ contains a subdiagram of the type F_4 or H_4 , and then finish the proof by really boring lemmas 6.6 – 6.9 concerning the subdiagrams F_4 and H_4 .

We say that a Coxeter diagram satisfies the *signature condition* if it is admissible and its positive inertia index does not exceed 7.

Recall that if $u, v \in \Sigma$ then $\langle u, v \rangle = m$ (∞ or 2) means that u and v are joined by an $(m - 2)$ -fold edge (dotted or empty edge, respectively).

Existence of a subdiagram of the type F_4 or H_4

In this subsection we show the following properties of Σ :

- Any node of Σ belongs to at most one dotted edge (Lemma 6.2);
- Σ contains no subdiagram of the type $G_2^{(k)}$ for $k > 5$ (Lemma 6.4);
- Σ contains at least one subdiagram of the type F_4 or H_4 (Lemma 6.5), and any such subdiagram has at least 2 bad neighbors (Lemma 6.3).

Recall from Lemma 1.3 that any elliptic subdiagram of Σ has at most 3 bad neighbors.

Lemma 6.1. *A subdiagram of the type $G_2^{(k)}$, $k > 3$, has at most 2 bad neighbors.*

Proof. Suppose that $S_0 \subset \Sigma$ is a subdiagram of the type $G_2^{(k)}$, $k > 3$, with 3 bad neighbors. Then $P(S_0)$ is a Coxeter 5-simplex, which is impossible. □

Lemma 6.2. *Any node of Σ is incident to at most one dotted edge.*

Proof. Suppose that a node v is incident to at least two dotted edges. Then the facet f of P corresponding to v is a (possibly non-Coxeter) 6-polytope with at most $6 + 2$ facets. By Cor. 3.1, f is not a simplex. Hence (by Prop. 1.7), it is a product of two simplices, i.e. either $\Delta^5 \times \Delta^1$, or $\Delta^4 \times \Delta^2$, or $\Delta^3 \times \Delta^3$. The first case is impossible, since Σ contains no large missing faces. The second and the third cases are impossible since the diagrams $[5, 3]_1$ and $[4, 4]_1$ have no 0- and 1-liftings with positive inertia index smaller than 8 (see Lemmas 3.6 and 3.11). □

Lemma 6.3. *Any subdiagram of Σ of the type H_4 or F_4 has at least two neighbors.*

Proof. Suppose that S_0 is a diagram of the type H_4 or F_4 . Since Σ is a connected diagram, S_0 has at least one neighbor. Suppose that S_0 has a unique neighbor, a . Then $P(S_0)$ is a Coxeter 3-polytope with $3 + 3$ facets. There are two simple combinatorial types of 3-polytopes with 6 facets, namely, a cube and a doubly truncated tetrahedron, i.e. a polytope with 2 pentagonal, 2 quadrilateral and 2 triangular facets. The later case is impossible for $P(S_0)$, since any of its triangular facets does not meet two other facets in contradiction to lemma 6.2 (here we use that $\Sigma_{S_0} = \overline{S_0} \subset \Sigma$, since S_0 is a diagram of the type F_4 or H_4). Hence, $P(S_0)$ is a cube. Denote by b_1 and b_2 , c_1 and c_2 , d_1 and d_2 the ends of the dotted edges in $\overline{S_0} = \Sigma_{S_0}$. By Lemma 1.4, a is joined with each of the dotted edges b_1b_2 , c_1c_2 and d_1d_2 . We assume that a is joined with b_1 , c_1 and d_1 .

Suppose that $[b_1, c_1] \geq 4$. Then the subdiagram $\langle b_1, c_1 \rangle$ has at least 3 bad neighbors (b_2 , c_2 and a), which is impossible by Lemma 6.1. Furthermore $[b_1, c_1] \neq \infty$, since $P(S_0)$ is a cube. Hence, $[b_1, c_1] = 2$ or 3. Similarly, $[b_1, d_1] \leq 3$ and $[c_1, d_1] \leq 3$.

Suppose that $[a, b_1] \geq 4$. Let $S_1 = \langle a, b_1 \rangle$. If $[a, b_1] \geq 6$, then S_1 has at least 3 bad neighbors b_2, c_1 and d_1 in contradiction to Lemma 6.1. Hence, $[a, b_1] = 4$ or 5 . By Lemma 6.1, the diagram S_1 has at most 2 bad neighbors, thus, at least one of c_1 and d_1 is a good neighbor of S_1 (recall that both c_1 and d_1 are joined with a). We may assume that c_1 is a good neighbor of S_1 and consider the diagram $S_2 = \langle S_1, c_1 \rangle$ of the type H_3 or B_3 . The diagram S_2 has at least four bad neighbors, namely b_2, c_2, d_1 and one of the nodes of S_0 (since a is a neighbor of S_0), which is impossible by Lemma 1.3. The contradiction shows that $[a, b_1] = 3$ ($[a, b_1] \neq \infty$ by Lemma 6.2). Similarly, $[a, c_1] = [a, d_1] = 3$.

Since Σ contains no parabolic subdiagrams and $[b_1, c_1] \leq 3$, we obtain that $[b_1, c_1] = 2$. Similarly, $[b_1, d_1] = [c_1, d_1] = 2$, so the diagram $\langle a, b_1, c_1, d_1 \rangle$ is a diagram of the type D_4 . This diagram has at least four bad neighbors, b_2, c_2, d_2 and one of x_i , $1 \leq i \leq 4$. We come to a contradiction, and the lemma is proved. $\square$

Lemma 6.4. Σ contains no subdiagram $G_2^{(k)}$ for $k > 5$.

Proof. Suppose that Σ contains a subdiagram $S_0 = \langle x_1, x_2 \rangle$ of the type $G_2^{(k)}$, $k > 5$. Without loss of generality we may assume that the edge of S_0 has the maximal multiplicity amongst all edges in Σ . Since Σ is connected, S_0 has at least one (evidently bad) neighbor; by Cor. 1.3, S_0 has at most 2 neighbors. Hence, S_0 has either 1 or 2 neighbors. We consider these two cases.

Case 1. Suppose that S_0 has a unique neighbor a . Then $P(S_0)$ is a 5-polytope with $5+3$ facets. Corollary 1.1 implies that $\Sigma_{S_0} = \overline{S}_0$. Hence, by Lemma 6.2, any node of Σ_{S_0} is incident to at most one dotted edge. The list of 5-polytopes with 8 facets contains a unique entry satisfying this condition (see Fig. 8(a) for this diagram and notation for its nodes). By Lemma 1.4, a is joined either with z_1 or with z_2 , say with z_1 . Denote $S_1 = \langle b_1, z_1 \rangle$. If a is a bad neighbor of S_1 , then S_1 has 3 bad neighbors (y_1, z_2, a) in contradiction to Lemma 6.1. Therefore, a is a good neighbor of S_1 , $[a, z_1] = 3$, and the diagram $S_2 = \langle a, S_1 \rangle$ is a diagram of the type B_3 . Recall that a is a neighbor of S_0 , and we may assume that x_1 is joined with a . If x_1 is a good neighbor of S_2 , then the diagram $\langle x_1, S_2 \rangle$ has more than 3 bad neighbors (namely, x_2, z_2, y_1 and some node of the Lannér diagram $\langle b_2, y_4, y_3, y_2 \rangle$ attached to a ; the latter neighbor is a bad one since a is not a leaf of $\langle x_1, S_2 \rangle$). This is impossible by Lemma 6.1, so x_1 is a bad neighbor of S_2 and S_2 has 3 bad neighbors (x_1, y_1, z_2) . Thus, $P(S_2)$ is a 4-dimensional simplex and Σ_{S_2} is a Lannér diagram of order 5. By Cor. 1.2, this implies that $\overline{S}_2$ is a Lannér diagram, too. However, $\overline{S}_2 = \langle x_2, b_2, y_4, y_3, y_2 \rangle$ cannot be a Lannér diagram, since it contains a Lannér diagram $\langle b_2, y_4, y_3, y_2 \rangle$ of order 4. The contradiction shows that S_0 has 2 neighbors.

Case 2. Suppose that the diagram $S_0 = \langle x_1, x_2 \rangle$ has 2 neighbors a_1 and a_2 . Then $P(S_0)$ is a 5-prism. We have two possibilities for this prism shown in Fig. 8(b),(c). We denote the nodes of $\overline{S}_0$ as it is shown in the figure. Denote by S_1 the subdiagram $\langle y_1, y_2, y_3, y_4 \rangle$ of the type F_4 or H_4 . Since the subdiagram $\overline{S}_1 = \Sigma_{S_1}$ contains the diagram S_0 of the type $G_2^{(k)}$, $k > 5$, the diagram $\overline{S}_1$ is not a Lannér diagram of order 4, so the face $P(S_1)$ is not a 3-simplex. Hence, S_1 has at most two neighbors, and at least one of a_1 and a_2 is not a neighbor of S_1 . We may assume that a_1 is not a neighbor of S_1 . By Lemma 1.4, this means that a_1 is joined with y_5 (since y_5 belongs to a Lannér diagram $\langle y_1, y_2, y_3, y_4, y_5 \rangle \subset \overline{S}_0$). We consider cases $S_1 = F_4$ and $S_1 = H_4$ separately.

In the case $S_1 = F_4$ consider two subdiagrams $\langle y_2, y_3, y_4, y_5, z_1 \rangle$ and $\langle y_3, y_2, y_1, y_5, z_1 \rangle$ of the type B_5 . Each of these diagrams has 3 bad neighbors (a_1, y_1, z_2 and a_1, y_4, z_2 respectively), so a_2 is not a bad neighbor for these diagrams. Therefore, a_2 is not joined with a Lannér diagram $\langle S_1, y_5 \rangle$, which contradicts Lemma 1.4.

Now we are left with the case $S_1 = H_4$. Since a_1 is not a neighbor of S_1 , the node a_2 is a neighbor of S_1 , otherwise S_1 has a unique neighbor in contradiction to Lemma 6.3. Therefore,




Figure 8: To the proof of Lemmas 6.4 and 6.6

S_1 has 2 bad neighbors, y_5 and a_2 , and $P(S_1)$ is a 3-prism. This means that the diagram $\overline{S}_1 = \Sigma_{S_1}$ consists of a dotted edge $z_1 z_2$ and a Lannér diagram of order 3 $\langle S_0, a_1 \rangle$. Thus, the subdiagram $X = \Sigma \setminus \langle z_1, z_2 \rangle = \langle S_0, a_1, S_1, y_5 \rangle$ consists of two Lannér diagrams $\langle S_0, a_1 \rangle$ and $\langle S_1, y_5 \rangle$ joined by a unique edge $a_1 y_5$. If this edge is not dotted one, then the subdiagram X is superhyperbolic, unless $[a_1, y_5] = 5$. We consider two cases, $[a_1, y_5] = 5$ and $[a_1, y_5] = \infty$.

Case 2.1. Suppose that $[a_1, y_5] = 5$. Then the subdiagram $\langle a_1, y_5, y_4, y_3 \rangle$ of the type H_4 has 3 bad neighbors, namely z_1, z_2 and one of x_1 and x_2 , say x_1 . Hence, z_2 is not a neighbor of $\langle a_1, y_5, y_4, y_3 \rangle$. In particular, $[z_2, a_1] = 2$, which means (Lemma 1.4) that $[a_1, z_1] \neq 2$. Therefore, z_1 is a bad neighbor of $S_2 = \langle a_1, y_5 \rangle$, and Σ_{S_2} is a diagram of a 5-polytope with at most 8 facets. Consider Σ_{S_2} . By Prop. 1.11, the subdiagram $S_1 = \langle y_1, y_2, y_3, y_4 \rangle$ turns into a linear diagram of order 4 with a triple edge $\tilde{y}_1 \tilde{y}_2$, simple edge $\tilde{y}_2 \tilde{y}_3$ and an edge $\tilde{y}_3 \tilde{y}_4$ labeled by 10 in Σ_{S_2} . However, no diagram of 5-polytope with at most 8 facets contains such a subdiagram. Thus, the case $[a_1, y_5] = 5$ is impossible.

Case 2.2. Suppose that $[a_1, y_5] = \infty$. Consider the subdiagram $S_2 = \langle y_1, y_2, y_3 \rangle$ of the type H_3 . If a_2 is a bad neighbor of S_2 then $P(S_2)$ is a 4-polytope with $4 + 3 = 7$ facets, and Σ_{S_2} contains a subdiagram $G_2^{(k)}$ for $k > 5$ and at least 2 dotted edges ($\tilde{z}_1 \tilde{z}_2$ and $\tilde{y}_4 \tilde{y}_5$). The list of 4-polytopes with 7 facets contains no entry with these properties. Hence, a_2 is not a bad neighbor of S_2 . If a_2 is a good neighbor of S_2 then the diagram $\langle y_1, y_2, y_3, a_2 \rangle$ of the type H_4 has at least 4 bad neighbors (namely, y_4 and at least one of x_1 and x_2 , one of z_1 and z_2 , and one of a_1 and y_5). Therefore, a_2 is not a neighbor of S_2 , and hence, by Lemma 6.3, a_2 is joined with y_4 . Consider the subdiagram $S_3 = \langle y_2, y_3, y_4, y_5, z_1 \rangle$ of the type A_5 or B_5 . S_3 has three bad neighbors (y_1, a_1, z_2), so a_2 is a good neighbor of S_3 and $\langle a_2, S_3 \rangle$ is a diagram of the type E_6 having at least four bad neighbors (y_1, z_2, a_1 and one of x_1 and x_2). This contradiction proves the lemma. $\square$

Lemma 6.5. Σ contains at least one subdiagram of the type F_4 or H_4 .


Proof. Suppose that the lemma is broken, i.e. Σ contains no subdiagram of the types F_4 and H_4 .

Suppose that Σ contains a subdiagram $S_0 = \langle x_1, x_2 \rangle$ of the type $G_2^{(4)}$ or $G_2^{(5)}$, having a bad neighbor. Then, $P(S_0)$ is a 5-polytope with at most $5 + 3$ facets. Therefore, Σ_{S_0} contains a subdiagram of the type either F_4 or H_4 . Cor. 1.2 implies that $\overline{S}_0$ contains a subdiagram of this type, which contradicts the assumption.

In particular, we conclude that Σ contains no Lannér diagram of order 3 (we use also Lemma 6.4). Since any Lannér diagram of order 5 contains a subdiagram of type F_4 or H_4 , Σ contains no Lannér diagram of order 5 either. By [E1, Satz 6.9], any simple d -polytope ($d > 4$) with $d + 4$ facets contains at least one missing face of order greater than 2. Thus, Σ contains a Lannér subdiagram L of order 4. Let $S_0 \subset L$ be a subdiagram of the type H_3 or B_3 . Denote $a_1 = L \setminus S_0$ and $S_0 = \langle x_1, x_2, x_3 \rangle$.


Suppose that S_0 has three bad neighbors. Then $P(S_0)$ is a 4-simplex, so Σ_{S_0} (and, hence, $\overline{S}_0$) contains a subdiagram of the type either F_4 or H_4 , which is impossible by the assumptions.

Suppose that S_0 has a unique bad neighbor, a_1 . Then $P(S_0)$ is a 4-polytope with $4 + 3$ facets. It follows from the assumption of the lemma that Σ_{S_0} is a diagram of a 4-polytope with $4 + 3$ facets containing neither a subdiagram of type F_4 nor a subdiagram of the type H_4 , and containing no subdiagram of the type $G_2^{(k)}$, $k > 5$ (Lemma 6.4). Each of these diagrams contains a subdiagram $S_1 = \langle y_1, y_2 \rangle$ of the type $G_2^{(4)}$ or $G_2^{(5)}$ having a bad neighbor. If S_0 is a diagram of the type H_3 , then it has no good neighbor (by the assumption Σ contains no subdiagram of the type H_4), and $\overline{S}_0 = \Sigma_{S_0}$ contains a subdiagram $\langle \tilde{y}_1, \tilde{y}_2 \rangle$ of the type $G_2^{(4)}$ or $G_2^{(5)}$ having a bad neighbor. As it is shown above, this is impossible. Thus, S_0 is a diagram of the type B_3 . By Cor. 1.1, either $S_1 = \langle y_1, y_2 \rangle \subset \overline{S}_0$ is a multiple edge with a bad neighbor (which is impossible) or we have one of the following possibilities for the diagram Σ_{S_0} :



In this case the double edge $\tilde{y}_1 \tilde{y}_2$ may turn into a simple edge $y_1 y_2$ in $\overline{S}_0$, that leads to a subdiagram $\langle y_1, y_2, y_3, y_4 \rangle \subset \overline{S}_0$ of the type H_4 , which contradicts the assumption. Therefore, the multiple edge $\tilde{y}_1 \tilde{y}_2$ remains multiple edge $y_1 y_2$ in $\overline{S}_1$, and Corollary 1.1 implies that the bad neighbor of S_1 remains bad in $\overline{S}_0$, which is impossible. So, S_0 cannot have 3 bad neighbors.

Therefore, S_0 has exactly 2 bad neighbors, a_1 and a_2 , and $P(S_0)$ is either an Esselmann polytope or a 4-prism. Since Σ contains no diagram of the types F_4 and H_4 , using Cor. 1.2 we obtain that Σ_{S_0} is the following diagram:



By Cor. 1.2, the nodes of Σ_{S_0} (with possible exclusion for y_6 in case of $[y_4, y_6] = 2$) cannot be good neighbors of S_0 . In particular, $\overline{S}_0$ contains a cyclic Lannér diagram of order 4 with a unique double edge, and contains no subdiagram of the type H_3 . Furthermore, it is easy to check that a dotted edge $\tilde{y}_5 \tilde{y}_6$ of Σ_{S_0} corresponds to a dotted edge $y_5 y_6$ in $\overline{S}_0$ (indeed, otherwise $[y_4, y_6] = 2$, and if the edge $y_5 y_6$ is triple one, then $\langle y_3, y_4, y_5, y_6 \rangle$ is a subdiagram of the type H_4 , which contradicts the assumption; if $y_5 y_6$ is a double edge, then $\langle y_2, y_3, y_4, y_5, y_6 \rangle$ is a parabolic subdiagram of Σ of the type $\tilde{C}_4$, which is also impossible; if $y_5 y_6$ is a simple edge, then $\langle y_5, y_6, S_0 \rangle$ is a diagram of the type B_5 and Prop. 1.11 shows that $\tilde{y}_5 \tilde{y}_6$ must not be a dotted but a double edge of Σ_{S_0}). So $\overline{S}_0$ consists of a dotted edge and of a cyclic Lannér diagram of order 4, and $\overline{S}_0 = \Sigma_{S_0}$.

Consider the diagram $S_1 = \langle y_1, y_2, y_3 \rangle$. It is a subdiagram of the type B_3 in the Lannér diagram $\langle y_1, y_2, y_3, y_4 \rangle$ of order 4. The same reasoning as for $\overline{S}_0$ shows that the diagram S_1 has exactly 2 bad neighbors (y_4 and one of a_1 and a_2), and the diagram $\overline{S}_1$ consists of a dotted edge $y_5 y_6$ and a cyclic Lannér diagram (either $\langle S_0, a_2 \rangle$ or $\langle S_0, a_1 \rangle$ respectively). Without loss of generality we may assume that a_2 is a bad neighbor of S_1 (and $\langle a_1, S_0 \rangle$ is a cyclic Lannér diagram). By Cor. 1.2, a_1 is not a good neighbor of S_1 , so a_1 is joined with y_4 (Lemma 1.4). If $[a_1, y_4] = 3$ or 4 then the subdiagram $\langle x_1, x_2, a_1, y_2, y_3, y_4 \rangle$ contains a parabolic subdiagram of the type $\tilde{B}_5$ or $\tilde{B}_3$. If $[a_1, y_4] = 5$ then $\langle x_2, x_3, a_1, y_4 \rangle$ is a subdiagram of the type H_4 . Therefore, $[a_1, y_4] = \infty$.

Consider the diagram $S_2 = \langle y_2, y_3, y_4 \rangle$. It has two bad neighbors, a_1 and y_1 , so a_2 is not a bad neighbor of S_2 . Therefore, a_2 is joined with y_1 , and repeating the reasoning above we obtain that $[a_2, y_1] = \infty$. Therefore, the diagram $\langle y_1, y_3, y_4, y_5 \rangle$ of the type D_4 has 4 bad neighbors a_1, a_2, y_2, y_6 , which is impossible. $\square$

Treatment of the subdiagrams F_4 and H_4

We have proved that Σ contains at least one subdiagram of the type F_4 or H_4 . In Lemmas 6.6 and 6.7 we prove that such a subdiagram always has 3 bad neighbors. Next, in Lemma 6.8 we show that Σ contains no subdiagram of the type F_4 . Lemma 6.9 finishes the proof of Theorem 3. Notice, that the proofs of Lemmas 6.7 and 6.9 (concerning the subdiagrams of the type H_4) turn out to be much more complicated than the proofs of the similar Lemmas 6.6 and 6.8 (concerning the subdiagrams of the type F_4). The possible reason is that a diagram of the type H_4 appears in diagrams of d -polytopes with at most $d + 3$ facets more often than a diagram of the type F_4 does.

Lemma 6.6. *Any subdiagram of Σ of the type F_4 has three neighbors.*

Proof. Suppose the contrary. By Lemma 6.3, this implies that Σ contains a subdiagram S_0 of the type F_4 with 2 neighbors. Then $P(S_0)$ is a 3-prism. The diagram $\Sigma_{S_0} = \overline{S}_0$ consists of a dotted edge $z_1 z_2$ and a Lannér diagram $\langle x_1, x_2, x_3 \rangle$ of order 3, which in its turn contains some multiple edge. Choose in $L = \langle x_1, x_2, x_3 \rangle$ an edge $S_1 = \langle x_1, x_2 \rangle$ of maximal possible multiplicity in L . The diagram S_1 has at least one bad neighbor, x_3 . By Lemma 6.1, S_1 has either 1 or 2 bad neighbors.

Suppose that S_1 has a unique bad neighbor x_3 . Then $P(S_1)$ is a 5-polytope with $5 + 3 = 8$ facets. The diagram S_0 of the type F_4 is not joined with S_1 . Hence, the diagram Σ_{S_1} contains a subdiagram of the type F_4 , and Σ_{S_1} is the diagram shown in Fig. 8(a). We denote the nodes of $S_0 \subset \overline{S}_1$ by y_1, y_2, y_3, y_4 and denote the neighbors of S_0 by b_1 and b_2 . By Cor. 1.2, the nodes b_1 and b_2 are not neighbors of S_1 . A Lannér diagram $\langle S_1, x_3 \rangle$ should be connected with a Lannér diagram $\langle b_1, y_1, y_2, y_3 \rangle$, thus, x_3 is joined with b_1 . Similarly, x_3 is joined with b_2 . Consider the diagram $S_2 = \langle b_1, y_1, y_2 \rangle$ of the type H_3 . It has 3 bad neighbors (y_3, x_3 and z_1) and has no good neighbor. Therefore, $P(S_2)$ is a 4-simplex and $\overline{S}_2 = \Sigma_{S_2} = \langle y_4, b_2, z_2, x_1, x_2 \rangle$ is a Lannér diagram of order 5. Since $y_4 b_2$ and $x_1 x_2$ are two unjoined multiple edges, the diagram $\overline{S}_2$ is a linear Lannér diagram (with edges labeled by 5, 3, 3, 4 or by 5, 3, 3, 5). Since $[y_4, b_2] = 5$, the diagram $\langle y_4, b_2, z_2, x_1 \rangle$ (or $\langle y_4, b_2, z_2, x_2 \rangle$) is a diagram of the type H_4 . In particular, $[b_2, z_2] = 3$ in Σ . So, a simple edge $b_2 z_2$ of Σ turns into a double edge $b_2 \tilde{z}_2$ in Σ_{S_1} . In view of Cor. 1.1, this implies that $[x_1, x_2] = 4$. Since the multiplicity of $x_1 x_2$ is maximal in the Lannér diagram $\langle x_1, x_2, x_3 \rangle$, we obtain that x_3 is joined with both x_1 and x_2 . Therefore, the diagram $\langle y_4, b_2, z_2, x_1 \rangle$ (or $\langle y_4, b_2, z_2, x_2 \rangle$) of the type H_4 has at least 4 neighbors: y_3, x_2 (or x_1), x_3 and z_1 , which is impossible.

The contradiction shows that S_1 has 2 bad neighbors, x_3 and some node a_1 . So, $P(S_1)$ is a 5-prism containing a subdiagram of the type F_4 (see Fig. 8(c) for the notation). Recall that $a_1 \notin \overline{S}_1$, and hence, a_1 is a bad neighbor of $S_0 = \langle y_1, y_2, y_3, y_4 \rangle$. Consider the diagrams $\langle y_2, y_3, y_4, y_5, z_1 \rangle$ and $\langle y_3, y_2, y_1, y_5, z_1 \rangle$ of the type B_5 . The node a_1 is a bad neighbor for at least one of these diagrams, say $\langle y_2, y_3, y_4, y_5, z_1 \rangle$. Then this diagram $\langle y_2, y_3, y_4, y_5, z_1 \rangle$ has at least 4 bad neighbors (y_1, a_1, z_2 and one of x_1, x_2, x_3). The contradiction proves the lemma. $\square$

Lemma 6.7. *Any subdiagram of Σ of the type H_4 has three neighbors.*

Proof. Suppose the contrary. By Lemma 6.3, this implies that Σ contains a subdiagram S_0 of the type H_4 with 2 neighbors. Then $P(S_0)$ is a 3-prism. The diagram $\Sigma_{S_0} = \overline{S}_0$ consists of a

dotted edge z_1z_2 and a Lannér diagram $L = \langle y_1, y_2, y_3 \rangle$ of order 3, which in its turn contains some multiple edge. We suppose that y_1y_2 is an edge of maximal possible multiplicity in L , and let $S_1 = \langle y_1, y_2 \rangle$. By Lemma 6.1, S_1 has either 1 or 2 neighbors.

Case 1. Suppose that S_1 has a unique bad neighbor y_3 . Then $P(S_1)$ is a 5-polytope with $5 + 3 = 8$ facets. Since Σ_{S_1} contains a subdiagram S_0 of the type H_4 , the diagram Σ_{S_1} is the diagram of one of three types (a), (b) and (c) shown in Fig. 9. We denote the nodes of $S_0 \subset \overline{S_1}$ by x_1, x_2, x_3, x_4 and denote the neighbors of S_0 by a_1 and a_2 (see Fig. 9). We consider cases (a), (b) and (c) separately.





Figure 9: Possibilities for Σ_{S_1} (Case 1).

Case 1.1. Suppose that Σ_{S_1} is the diagram shown in Fig. 9(a). Recall that z_1z_2 is a dotted edge in Σ , so Lemma 6.2 implies that a_1 is a good neighbor of S_1 . Without loss of generality we may assume that $[a_1, y_1] = 3$ and $[a_1, y_2] = 2$. On the other hand, a_2 is not a good neighbor of $S_1 = \langle y_1, y_2 \rangle$, since $[a_2, z_2] = 5$. Hence, a_2 is joined with y_3 (otherwise a Lannér diagram $\langle x_1, x_2, x_3, x_4, a_2 \rangle$ is not joined with a Lannér diagram $L = \langle S_1, y_3 \rangle$). Notice that the diagram $S_2 = \langle z_2, a_2, x_4, x_3 \rangle$ of the type H_4 has 3 neighbors (x_2, z_1, y_3) , hence, $P(S_2)$ is a 3-simplex, so the diagram $\tilde{S}_2 = \langle x_1, a_1, y_1, y_2 \rangle$ is a Lannér diagram. Furthermore, $\tilde{S}_2$ is a linear Lannér diagram (x_1 is not joined with $\langle y_1, y_2 \rangle$, and a_1 is not joined with y_2 and joined with y_1 by a simple edge). Therefore, $[x_1, a_1] = 4$ or 5 . Consider the diagram $S_3 = \langle x_1, a_1, y_1 \rangle$ of the type H_3 or B_3 . It has 3 bad neighbors (x_2, z_1, y_2) , hence $P(S_3)$ is a 4-simplex, and Σ_{S_3} is a Lannér diagram of order 5. By Cor. 1.2, this implies that $\tilde{S}_0$ is a Lannér diagram, too. However, in $\tilde{S}_3 = \langle x_3, x_4, a_2, z_2, y_3 \rangle$ one end (a_2) of the triple edge has valency at least three, which is impossible in a Lannér diagram of order 5.

Case 1.2. Suppose that Σ_{S_1} is one of the diagrams shown in Fig. 9(b) and (c). In these cases neither a_1 nor a_2 can be a good neighbor of S_1 , and Lemma 6.2 implies that both z_1 and z_2 are good neighbors of S_1 (recall that z_1z_2 is a dotted edge in Σ). By Lemma 6.4, $[a_i, z_i] = 3, 4$ or 5 ($i = 1, 2$). If $[a_1, z_1] = 5$, then the diagram $\langle z_1, a_1, x_4, x_3 \rangle$ of the type H_4 has at least four neighbors $(x_2, z_2, a_2$ and at least one of y_1 and y_2). If $[a_1, z_1] = 4$, then the diagram $\langle z_1, a_1, x_4, x_3, x_2 \rangle$ of the type B_5 has at least four bad neighbors $(x_1, z_2, a_2$ and at least one of y_1 and y_2). Finally, if $[a_1, z_1] = 3$, then the edge $\tilde{a}_1\tilde{z}_1$ of Σ_{S_1} should be either a double edge or an edge marked by 10, but not a dotted edge.

Case 2. Suppose that S_1 has 2 bad neighbors: y_3 and one of a_1 and a_2 , say a_1 . Then, $P(S_1)$ is a 5-prism containing a subdiagram of the type H_4 , so, the diagram Σ_{S_1} is the following one:



By Cor. 1.2, a_2 is not a good neighbor of S_1 , hence, a_2 is a neighbor of y_3 (see Lemma 1.4). If $[a_2, y_3] \neq \infty$, then the subdiagram $\langle S_0, a_2, y_3, S_1 \rangle$ is superhyperbolic (in assumption that Σ contains no subdiagrams of the type $G_2^{(k)}$ for $k > 5$ and the multiplicity of the edge $y_1 y_2$ is maximal possible in $\langle y_1, y_2, y_3 \rangle$). Hence, we assume $[a_2, y_3] = \infty$. Let $S_4 = \langle x_1, x_2, x_3 \rangle$. Consider 3 cases: a_1 is either a bad neighbor of S_2 , or a good neighbor, or a non-neighbor.

Case 2.1. Suppose that a_1 is a bad neighbor of S_4 . Then $P(S_4)$ is a 4-polytope with $4 + 3$ facets, and Σ_{S_4} contains at least 3 dotted edges $\tilde{z}_1 \tilde{z}_2$, $\tilde{a}_2 \tilde{y}_3$ and $\tilde{x}_4 \tilde{a}_2$. However, no Coxeter diagram of 4-polytope with $4 + 3$ facets contains more than 3 dotted edges and if a Coxeter diagram of a 4-polytope with $4 + 3$ facets have exactly 3 dotted edges then any dotted edge is incident to some other dotted edge. The edge $\tilde{z}_1 \tilde{z}_2$ has no common vertices with $\tilde{a}_2 \tilde{y}_3$ and $\tilde{x}_4 \tilde{a}_2$, so we obtain a contradiction.

Case 2.2. Suppose that a_1 is a good neighbor of $S_4 = \langle x_1, x_2, x_3 \rangle$. Consider the diagram $S_5 = \langle x_1, x_2, x_3, a_2 \rangle$ of the type H_4 . S_5 has 3 bad neighbors: x_4 , one of y_1 and y_2 (say y_1), and one of z_1 and z_2 . This implies that y_2 is not a neighbor of S_5 , in particular, $[a_1, y_2] = 2$ and $[a_1, x_1] = 2$. On the other hand, a_1 is a bad neighbor of $\langle y_1, y_2 \rangle$. Therefore, $[a_1, y_1] \in \{4, 5, \infty\}$.

Consider the subdiagram $S_6 = \langle x_2, x_3, x_4, a_2, z_1 \rangle$ of the type either A_5 or B_5 . Since a_1 is a neighbor of $S_0 = \langle x_1, x_2, x_3, x_4 \rangle$ and as we have shown above, $[a_1, x_1] = 2$, we obtain that a_1 is a neighbor of S_6 . The diagram S_6 has 3 bad neighbors: x_1, z_2, y_3 . Hence, a_1 is a good neighbor of S_6 . Therefore, $\langle a_1, S_6 \rangle$ is a diagram of the type D_6 with 3 bad neighbors (x_1, z_2, y_3). However, since $[a_1, y_1] = 4, 5$ or ∞ , y_1 is also a bad neighbor of this diagram, which is impossible.

Case 2.3. Suppose that a_1 is not a neighbor of $S_4 = \langle x_1, x_2, x_3 \rangle$. Then a_1 is joined with x_4 (since a_1 is a neighbor of $S_0 = \langle x_1, x_2, x_3, x_4 \rangle$). Consider the subdiagram $S_7 = \langle x_2, x_3, x_4, a_2, z_1 \rangle$ of the type A_5 or B_5 . S_7 has 3 bad neighbors: x_1, z_2, y_3 . Thus, a_1 is a good neighbor of S_7 , and $\langle S_7, a_1 \rangle$ is a diagram of the type E_6 . However, this diagram $\langle S_7, a_1 \rangle$ has 4 bad neighbors: x_1, z_2, y_3 and one of y_1 and y_2 (since a_1 is a bad neighbor of S_1 by assumption).

This shows that the assumption that S_1 has 2 bad neighbors is impossible, which completes the proof of the lemma. $\square$

Lemma 6.8. Σ contains no subdiagram of the type F_4 .

Proof. Suppose that $S_0 = \langle x_1, x_2, x_3, x_4 \rangle$ is a subdiagram of Σ of the type F_4 . By Lemma 6.6, S_0 has 3 neighbors a_1, a_2 and a_3 . Then $P(S_0)$ is a 3-simplex, and $\overline{S}_0 = \Sigma_{S_0} = \langle y_1, y_2, y_3, y_4 \rangle$ is a Lannér diagram of order 4. Let $S_1 = \langle y_1, y_2, y_3 \rangle$ be a subdiagram of $\overline{S}_0$ of the type H_3 or B_3 . S_1 has at least one bad neighbor, y_4 . Let S_2 be a subdiagram of $\overline{S}_0$ of the type H_3 or B_3 containing y_4 (it does exist by Lemma 1.5), we may assume that $S_2 = \langle y_2, y_3, y_4 \rangle$.

The diagram S_1 has at least one bad neighbor, y_4 . We consider three cases where S_1 has 1, 2 or 3 bad neighbors respectively.

Case 1. Suppose that S_1 has a unique bad neighbor y_4 . Then $P(S_1)$ is a 4-polytope with $4 + 3$ facets.

Suppose in addition that the diagram $\langle a_i, S_0, \overline{S}_0 \rangle$ contains a dotted edge for any $i = 1, 2, 3$. Then each of a_i ($i = 1, 2, 3$) is an end of some dotted edge. Consider three other ends. If S_0 contains three ends of dotted edges, then Σ_{S_1} contains three dotted edges with mutually distinct ends (since $S_0 \subset \Sigma_{S_1}$ and $a_i \in \Sigma_{S_1}$ for $i = 1, 2, 3$). This is impossible for a diagram of a

4-polytope with $4 + 3$ facets. Therefore, at least one of the ends of the dotted edges belongs to $\overline{S}_0$. Since $a_i \in \Sigma_{S_1}$ ($i = 1, 2, 3$), S_1 contains no end of a dotted edge. Hence, y_4 is the only end of a dotted edge in $\overline{S}_0$. Then S_2 has at least 2 bad neighbors (namely, y_1 and some a_j such that $[a_j, y_4] = \infty$). If S_2 has exactly 2 bad neighbors, then Σ_{S_2} is a diagram of either an Esselmann polytope or of a 4-prism. However, it contains two dotted edges, which is impossible. If S_2 has 3 bad neighbors, then Σ_{S_2} is a diagram of a 4-simplex. At the same time, it contains at least one dotted edge.

Therefore, we may assume that the diagram $\langle a_1, S_0, \overline{S}_0 \rangle$ contains no dotted edges. In view of Lemma 6.4, we are left with finitely many possibilities for the diagram $\langle a_1, S_0, \overline{S}_0 \rangle$. The only two of these diagrams satisfying the signature condition and the condition that S_1 has a unique bad neighbor are the diagrams shown in Fig. 10.

For the diagram shown in Fig. 10(a) consider the subdiagram $S_3 = \langle a_1, y_4 \rangle$ having 2 bad neighbors. Then Σ_{S_3} is a diagram of a 5-prism, but Σ_{S_1} contains a diagram $\langle \tilde{x}_1, \tilde{x}_2, \tilde{x}_3, \tilde{y}_1, \tilde{y}_2 \rangle$ of the type $B_3 + B_2$ which is impossible for a diagram of a 5-prism.

Now, consider the diagram shown in Fig. 10(b). The subdiagram $S_4 = \langle x_4, a_1, y_3, y_2 \rangle$ of the type H_4 has 3 bad neighbors x_3, y_4 and y_1 . Therefore, a_2 and a_3 are not neighbors of S_4 . In particular, $[a_i, y_3] = [a_i, y_2] = 2$ for $i = 2, 3$.

Suppose that $[a_2, y_4] \neq 0$. Then the diagram $S_5 = \langle a_1, y_3, y_4 \rangle$ of the type H_3 has 3 bad neighbors (x_4, y_2 and a_2). Hence, $[a_3, y_4] = 2$, and $[a_3, y_1] \neq 0$ (see Lemma 1.4 applied to a Lannér diagram $L = \langle y_1, y_2, y_3, y_4 \rangle$). Furthermore, $P(S_5)$ is a simplex and Σ_{S_5} is a Lannér diagram of order 5. By Cor. 1.2, $\overline{S}_5$ is also a Lannér diagram. Since $\overline{S}_5$ contains the diagram S_0 of the type F_4 , $\overline{S}_0$ is a cyclic Lannér diagram and $[a_3, x_1] = [a_3, x_4] = 3$. However, the diagram $\langle x_1, a_3, x_4, a_1 \rangle$ of the type H_4 has at least 4 neighbors (x_2, x_3, y_3 and y_1 , which is a neighbor of a_3). The contradiction shows that $[a_2, y_4] = 0$. Similarly, $[a_3, y_4] = 0$.

By Lemma 1.4, a_2 and a_3 should be joined with a Lannér diagram $\langle y_1, y_2, y_3, y_4 \rangle$, so, both a_2 and a_3 are joined with y_1 . Therefore, the diagram $\langle a_1, y_3, y_2, y_1 \rangle$ of the type B_4 has 4 bad neighbors a_2, a_3, x_4, y_4 , which is impossible.




Figure 10: Case 1: two possibilities for the diagram $\langle a_1, S_0, \overline{S}_0 \rangle$

Case 2. Suppose that S_1 has two bad neighbors, y_4 and a_1 . Then $P(S_1)$ is either an Esselmann polytope or a 4-prism. Notice that Σ_{S_1} contains a subdiagram S_0 of the type F_4 and contains no $G_2^{(k)}$ for $k > 5$ (see Cor. 1.1). There are two prisms and one Esselmann polytope satisfying these conditions (see Fig. 11). We consider these polytopes separately.

Case 2.1. Suppose that Σ_{S_1} is the diagram shown in Fig. 11(a). Then $\Sigma_{S_1} = \overline{S}_1$. The node a_1 is a neighbor of $S_0 = \langle x_1, x_2, x_3, x_4 \rangle$. Without loss of generality we may assume that a_1 is a neighbor of $\langle x_1, x_2, x_3 \rangle$. Then the subdiagram $\langle x_1, x_2, x_3, a_3 \rangle$ of the type F_4 has 4 bad neighbors (x_4, a_1, a_2 and some node of $\overline{S}_0$ joined with a_3), which is impossible.

Case 2.2. Suppose that Σ_{S_1} is the diagram shown in Fig. 11(b). Then again $\Sigma_{S_1} = \overline{S}_1$. In particular, a_2 and a_3 are non-neighbors of S_1 , and a_2 and a_3 are joined with y_4 . Furthermore, the diagram $\langle x_1, x_2, x_3, a_3 \rangle$ of the type F_4 has 3 neighbors: x_4, a_2 and y_4 . Therefore, a_1 is not a neighbor of this diagram, hence $[a_1, x_4] \neq 2$. Consider the diagram $S_6 = \langle a_2, x_1, x_2 \rangle$ of the type H_3 . Clearly, it has no good neighbors, and $\overline{S}_6 = \Sigma_{S_6}$. Furthermore, $\overline{S}_6$ contains a dotted




Figure 11: Diagrams of 4-polytopes with 6 facets containing F_4 and containing no $G_2^{(k)}$ for $k > 5$.

edge a_3x_4 , so, S_6 has at most 2 bad neighbors. Thus, it has exactly 2 bad neighbors, x_3 and y_4 . Therefore, the subdiagram $\langle a_3, x_4, a_1 \rangle$ belongs to the subdiagram $\overline{S}_6 = \Sigma_{S_6}$, which is a diagram of a 4-prism. Hence, the edge $\tilde{x}_4\tilde{a}_1$ is adjacent to a dotted edge in the diagram of a 4-prism. This implies that $[x_4, a_1] = 3$, and $\langle x_1, x_2, x_3, x_4, a_1 \rangle$ is a parabolic diagram of the type $\tilde{F}_4$, which is impossible.

Case 2.3. Suppose that Σ_{S_1} is the diagram shown in Fig. 11(c). We consider 2 cases: either $[a_3, x_4] = \infty$ or $[a_3, x_4] \neq \infty$ in Σ .


Case 2.3.1. Suppose that $[a_3, x_4] = \infty$. Since a_2 is not a good neighbor of S_1 , y_4 is the only node of $\overline{S}_0$ joined with a_2 .

Consider the diagram $S_7 = \langle a_2, x_1 \rangle$ of the type $G_2^{(5)}$. If S_7 has a bad neighbor, then $P(S_7)$ is a 5-polytope with at most $5 + 3$ facets. At the same time, Σ_{S_7} contains a subdiagram $\langle \tilde{x}_2, \tilde{x}_3, \tilde{x}_4, \tilde{a}_3 \rangle$, and $\tilde{x}_2\tilde{x}_3$ is a dotted edge in Σ_{S_7} . However, no diagram of a 5-polytope with at most $5 + 3$ facets contains two dotted edges ($\tilde{x}_2\tilde{x}_3$ and $\tilde{x}_4\tilde{a}_3$) joined by a unique simple edge ($\tilde{x}_3\tilde{x}_4$). Therefore, S_7 has no bad neighbors. In particular, $[a_2, y_4] = 3$.

Consider the diagram $\langle a_2, S_0, \overline{S}_0 \rangle$. It contains no dotted edges and we completely know this diagram modulo finitely many possibilities for a Lannér subdiagram $\overline{S}_0$. However, for any of these possibilities the diagram $\langle a_2, S_0, \overline{S}_0 \rangle$ does not satisfy the signature condition. Hence, the case $[a_3, x_4] = \infty$ is impossible.

Case 2.3.2 Suppose that $[a_3, x_4] \neq \infty$. Then a_3 is a good neighbor of S_1 and $[a_3, x_4] = 5$ (otherwise we have either $[a_3, x_4] = 3$ and $\langle a_3, S_0 \rangle$ is a parabolic diagram of the type $\tilde{F}_4$, or $[a_3, x_4] = 4$ and $\langle a_3, x_4, x_3, x_2 \rangle$ is a parabolic diagram of the type $\tilde{C}_4$).

Consider the subdiagram $X = \langle S_0, \overline{S}_0, a_3 \rangle$. We know this diagram modulo finitely many possibilities for $\overline{S}_0$ (S_0 is a diagram of the type F_4 , $\overline{S}_0$ is a Lannér diagram of order 4, $S_1 \subset \overline{S}_0$ is a diagram of the type H_3 or B_3 , a_3 is a good neighbor of S_1 , and a_3 is joined with S_0 as shown in Fig. 11(c)). The only diagram satisfying these conditions and the signature condition is the following:



Clearly, a_2 is not a good neighbor of S_1 . Hence, it is not a neighbor of $S_1 = \langle y_1, y_2, y_3 \rangle$ and $[a_2, y_4] \neq 2$ (Lemma 1.4). Therefore, the diagram $S_2 = \langle y_2, y_3, y_4 \rangle$ has 3 bad neighbors a_2, a_3, y_1 . Thus, Σ_{S_2} is a Lannér diagram of order 5 containing a subdiagram of the type F_4 . This implies that a_1 is joined with x_1 and x_4 . However, in this case the diagram $\langle x_4, a_3, y_3, y_2 \rangle$ of the type H_4 has 4 bad neighbors a_1, x_3, y_1, y_4 .

We considered two prisms and an Esselmann polytope and no possibilities were found, so the proof of Case 2 is completed.

Case 3. Suppose that S_1 has three bad neighbors, y_4 , a_1 and a_2 . Then $\overline{S}_1$ is a cyclic Lannér diagram of order 5, i.e. $\overline{S}_1 = \mathcal{L}_5^5$, and $[a_3, x_1] = [a_3, x_4] = 3$, $[a_3, x_2] = [a_3, x_3] = 2$. By Cor. 1.2, a_3 is not a good neighbor of S_1 , and y_4 is the only node of $\overline{S}_0$ joined with a_3 .

As it is shown in Case 1 and Case 2, we may assume that the diagram $S_2 = \langle y_2, y_3, y_4 \rangle$ has also 3 bad neighbors. The reasoning above shows that each of a_i , $i = 1, 2, 3$, is either a bad neighbor of S_2 or a non-neighbor, and exactly one of a_i is a non-neighbor. Thus, a_3 and one of a_1 and a_2 (say a_1) are bad neighbors of S_2 and a_2 is a non-neighbor. Moreover, $\langle a_1, S_0 \rangle$ is a cyclic Lannér diagram, and y_1 is the only node of $\overline{S}_0$ joined with a_1 . Without loss of generality we may assume that a_2 is a neighbor of $\langle x_1, x_2, x_3 \rangle$. Then the diagram $\langle a_1, x_1, x_2, x_3 \rangle$ of the type B_4 has 3 bad neighbors x_1, a_2 and a_3 . Hence, y_1 is a good neighbor of $\langle a_1, x_1, x_2, x_3 \rangle$. So, $[y_1, a_1] = 3$. Recall, that a_1 is a bad neighbor of S_1 . This implies that either $[y_1, y_2] = 4, 5$ or $[y_1, y_3] = 4, 5$. Therefore, either y_2 or y_3 is a bad neighbor of $\langle y_1, a_1, x_1, x_2, x_3 \rangle$, which is impossible. $\square$

Lemma 6.9. *If Σ contains a subdiagram of the type H_4 then Σ is a diagram Σ_{P_7} .*

Proof. Suppose that $S_0 = \langle x_1, x_2, x_3, x_4 \rangle$ is a subdiagram of Σ of the type H_4 . By Lemmas 6.3 and 6.7, S_0 has 3 neighbors a_1, a_2 and a_3 . Then $P(S_0)$ is a 3-simplex, and $\overline{S}_0 = \Sigma_{S_0} = \langle y_1, y_2, y_3, y_4 \rangle$ is a Lannér diagram of order 4. Let $S_1 = \langle y_1, y_2, y_3 \rangle$ be a subdiagram of $\overline{S}_0$ of the type H_3 or B_3 . S_1 has at least one bad neighbor, y_4 . We consider three cases in which S_1 has 1, 2 and 3 bad neighbors respectively.

Case 1. Suppose that S_1 has a unique bad neighbor y_4 . Then $P(S_1)$ is a 4-polytope with $4 + 3$ facets. The same reasoning as in Case 1 of Lemma 6.8 shows that for some $i \in \{1, 2, 3\}$ the diagram $\langle a_i, S_0, \overline{S}_0 \rangle$ contains no dotted edges. We assume that the diagram $\langle a_1, S_0, \overline{S}_0 \rangle$ contains no dotted edges. In view of Lemma 6.4, we are left with finitely many possibilities for the diagram $\langle a_1, S_0, \overline{S}_0 \rangle$. Recall that $S_1 \subset \overline{S}_0$ is a subdiagram of the type H_3 or B_3 with a unique bad neighbor. There are only three possibilities for the diagram $\langle a_1, S_0, \overline{S}_0 \rangle$ satisfying this condition together with the signature condition, namely, the diagrams shown in Fig. 12. We consider these diagrams separately.




Figure 12: Three possibilities for the diagram $\langle a_1, S_0, \overline{S}_0 \rangle$.

Consider the diagrams shown in Fig. 12(a) and 12(b). Clearly, $S_1 = \langle y_2, y_3, y_4 \rangle$. Suppose that a_2 is joined with y_3 or y_4 . Since a_2 is not a bad neighbor of S_1 , we obtain a subdiagram $\langle a_2, S_1 \rangle$ of the type F_4 , which contradicts Lemma 6.8. Hence, both a_2 and a_3 are joined with either y_1 or y_2 (Lemma 1.4), and the subdiagram $\langle y_1, y_2, y_3 \rangle$ of the type H_3 has 4 bad neighbors (y_4, a_1, a_2, a_3) , which is impossible.

Consider the diagram shown in Fig. 12(c). Without loss of generality we assume that $S_1 = \langle y_1, y_2, y_3 \rangle$. Each of the diagrams $S_2 = \langle a_1, y_1, y_2, y_3 \rangle$ and $S_3 = \langle a_1, y_4, y_3, y_2 \rangle$ of the

type H_4 has bad neighbors x_4 and y_1 (or y_4 respectively). Thus, each of these diagrams has at most one extra neighbor (Lemma 6.7). On the other hand, by Lemma 1.4, each of the nodes a_2 and a_3 is joined with the diagram $\langle a_1, y_1, y_2, y_3, y_4 \rangle$. Thus, we may assume that a_2 is not joined with S_2 and joined with y_4 , and a_3 is not joined with S_3 and joined with y_1 . Since S_1 has no bad neighbor besides y_4 , $[a_4, y_1] = 3$. Furthermore, $[a_3, a_1] = 2$ (otherwise the diagram $\langle S_1, a_3 \rangle$ of the type H_4 has 4 bad neighbors y_4, a_2, a_1 and some x_i , $i \in \{1, 2, 3, 4\}$). If $[a_1, x_4] = 2$, then $\langle x_4, a_1, y_4, y_1, y_2, a_3 \rangle$ is a parabolic diagram of the type $\tilde{D}_5$, hence, $[a_3, x_4] \neq 2$. Moreover, $[a_1, x_4] \neq 3$ (otherwise, $\langle x_4, a_1, y_1, a_3 \rangle$ is a parabolic diagram $\tilde{A}_3$). Therefore, the diagram $S_4 = \langle x_2, x_3, x_4, a_1, y_4, y_3 \rangle$ of the type A_6 has three bad neighbors x_1, y_2 and a_3 . Hence, a_2 is a good neighbor of S_4 , and $[a_2, y_4] = 3$, $[a_2, x_4] = [a_2, x_3] = [a_2, x_2] = 2$. Therefore, $\langle x_3, x_4, a_1, y_4, a_2, y_1, y_2 \rangle$ is a parabolic diagram of the type $\tilde{E}_6$, which is impossible.

Case 2. Suppose that S_1 has two bad neighbors, y_4 and a_1 . Then $P(S_1)$ is either an Esselmann polytope or a 4-prism, $\Sigma_{S_1} = \langle \tilde{x}_1, \tilde{x}_2, \tilde{x}_3, \tilde{x}_4, \tilde{a}_2, \tilde{a}_3 \rangle$.

A Coxeter diagram of an Esselmann polytope containing no subdiagram of the types F_4 and $G_2^{(k)}$, $k \geq 6$, is one of the two diagrams shown in Fig. 13. A Coxeter diagram of a 4-prism containing no subdiagram of the type F_4 is one of the diagrams shown in Fig. 14. The diagram (a) is drawn for 3 times, since there are 3 different possibilities for the location of S_0 in this diagram (in any other diagram containing two subdiagrams of the type H_4 these subdiagrams are permuted by an automorphism of the diagram).




Figure 13: Diagrams of Esselmann polytopes containing H_4 and containing no $G_2^{(k)}$ for $k > 5$.

We assume that $\langle \tilde{x}_1, \tilde{x}_2, \tilde{x}_3 \rangle$ is the subdiagram of Σ_{S_1} of the type H_3 . It turns out that in any possible case for the diagram $\overline{S}_1$ the subdiagram $\langle \tilde{x}_1, \tilde{x}_2, \tilde{x}_3 \rangle$ has a unique bad neighbor in Σ_{S_1} . We assume that $\tilde{a}_2$ is the bad neighbor of the diagram $\langle \tilde{x}_1, \tilde{x}_2, \tilde{x}_3 \rangle$, then $\tilde{a}_3$ is the remaining node of Σ_{S_1} .

First we consider the Esselmann polytopes and then prisms.

Case 2.1. Suppose that Σ_{S_1} is an Esselmann diagram. The subdiagram $S_5 = \langle x_1, x_2, x_3, a_3 \rangle$ of the type H_4 has 3 bad neighbors in Σ (a_2, x_4 and some node of $\overline{S}_0$ joined with a_3). Hence, a_1 is not a neighbor of $S_5 = \langle x_1, x_2, x_3, a_3 \rangle$, so a_1 is joined with x_4 (as a neighbor of S_0). Furthermore, the subdiagram $\langle S_0, a_3, \overline{S}_0 \rangle$ consists of two diagrams $\langle S_0, a_3 \rangle$ and $\overline{S}_0$ joined by the edge $a_3 y_4$ only (and this edge does exist by Lemma 1.4). Moreover, this edge is a dotted one, otherwise the diagram $\langle S_0, a_3, \overline{S}_0 \rangle$ is superhyperbolic. In particular, three bad neighbors of the diagram $S_5 = \langle x_1, x_2, x_3, a_3 \rangle$ are a_2, x_4, y_4 , so $\langle S_1, a_1 \rangle = \overline{S}_5 = \Sigma_{S_5}$ is a Lannér diagram of order 4.

Consider the diagram $S_6 = \langle a_3, x_4 \rangle$ of the type $G_2^{(5)}$. S_6 has at least two bad neighbors, y_4 and x_4 , and $P(S_6)$ is a 5-dimensional polytope with at most $5 + 2$ facets. This implies that $P(S_6)$ is a 5-prism. Notice that the subdiagram $X = \langle x_1, x_2, a_2, y_1, y_2, y_3 \rangle$ is not joined with S_6 in Σ , so it does not differ from the subdiagram $\langle \tilde{x}_1, \tilde{x}_2, \tilde{a}_2, \tilde{y}_1, \tilde{y}_2, \tilde{y}_3 \rangle$ of Σ_{S_6} . It is clear, that X does not contain dotted edges (a_2 cannot be joined with S_1 by a dotted edge, since a_2 is not a bad neighbor of S_1). Moreover, as a diagram of a 5-prism should not contain Lannér

diagrams of order 3, we obtain that a_2 is a good neighbor of S_1 , the diagram Σ_{S_1} is the one shown in Fig. 13(b), $[a_2, x_2] = 3$, and S_1 is a diagram of the type B_3 (if it is of the type H_3 , then $\tilde{a}_2\tilde{x}_2$ is a dotted edge in Σ_{S_6} , so it cannot be adjacent to a triple edge $\tilde{x}_1\tilde{x}_2$ in a diagram of a 5-prism). Without loss of generality we may assume that $[y_1, y_2] = 3$, $[y_2, y_3] = 4$. Then we obtain a linear diagram $\langle x_1, x_2, a_2, y_1, y_2, y_3 \rangle$ with the following labels on its edges: 5, 3, 3, 3, 4. This implies that $[a_1, y_3] = \infty$ in Σ_{S_6} , thus, $[a_1, y_3] \neq 2, 3$ in Σ (if $[a_1, y_3] = 3$, the edge $\tilde{a}_1\tilde{y}_3$ of Σ_{S_6} is labeled by 10, and it is not a dotted edge as it should be). However, the multiple edges a_1y_3 and y_3y_2 cannot be adjacent in Lannér diagram $\langle S_1, a_1 \rangle$ of order 4.

The contradiction shows that Σ_{S_1} is not a diagram of an Esselmann polytope.

Case 2.2. Suppose that Σ_{S_1} is a 4-prism. Notice, that in all diagrams of 4-prisms all edges incident to a_2 are simple, and by Cor. 1.2, a_2 cannot be a good neighbor of S_1 . Hence, a_2 is a neighbor of y_4 (Lemma 1.4).

Furthermore, suppose that $[a_2, y_4] \neq \infty$. Then the diagram $\langle S_0, a_2, \bar{S}_0 \rangle$ satisfies the following properties: it contains no dotted edges; y_4 is the only node of $\bar{S}_0$ joined with a_2 ; a_2 is a bad neighbor of $H_3 \subset S_0$ and a_2 is joined with S_0 by simple edges only. It is easy to check that no diagram satisfies these properties together with signature condition. We obtain that $[a_2, y_4] = \infty$.

Consider the diagrams of 4-prisms case-by-case.




Figure 14: Diagrams of 4-prisms containing H_4 .


Case 2.2.1. Suppose that Σ_{S_1} is the diagram shown in Fig. 14(a1). If a_1 is a neighbor of $\langle x_1, x_2, x_4 \rangle$ then $\langle x_2, x_1, a_2, x_4 \rangle$ is a diagram of the type H_4 with 4 neighbors x_3, a_3, a_1, y_4 (recall that $[a_2, y_4] = \infty$). If a_1 is not a neighbor of $\langle x_1, x_2, x_4 \rangle$, then $[a_1, x_3] \neq 2$ and $\langle x_1, x_2, x_3, a_3 \rangle$ is a diagram of the type H_4 with 4 neighbors $(x_3, a_2, a_1$ and some $y_i, i \in \{1, 2, 3, 4\}$, see Lemma 1.4).

Case 2.2.2. Suppose that Σ_{S_1} is one of the diagrams shown in Fig. 14(a2)-(c). Since the diagram $\langle x_2, x_1, a_2, a_3 \rangle$ of the type H_4 has 3 bad neighbors x_3, x_4 and y_4 (since $[a_2, y_4] = \infty$), the diagram $\langle S_1, a_1 \rangle$ is a Lannér diagram. Clearly, the nodes x_1, x_2, a_2 are not attached to this Lannér diagram (otherwise $\langle x_2, x_1, a_2, a_3 \rangle$ has 4 bad neighbors, x_3, x_4, y_4 and some node of $\langle S_1, a_1 \rangle$). Since a Lannér diagram $\langle a_1, S_1 \rangle$ is joined with a Lannér diagram $\langle x_1, x_2, x_3, a_2 \rangle$, we obtain that $[x_3, a_1] \neq 2$. The signature condition applied to the diagram $\langle S_0, a_1, a_2, S_1 \rangle$ implies that $[x_3, a_1] = \infty$. However, this means that the diagram $\langle x_2, x_3, x_4, a_2 \rangle$ of the type A_4 (in case (a2)), or $A_3 + A_1$ (in cases (a3) and (c)), or D_4 (in case (b)) has 4 bad neighbors, x_1, a_3, y_4, a_1 (again, we use that $[a_2, y_4] = \infty$).

Case 2.2.3. Suppose that Σ_{S_1} is the diagram shown in Fig. 14(d). Since the diagram $\langle x_2, x_1, a_2, x_4 \rangle$ of the type H_4 has 3 bad neighbors, x_3, a_3 and y_4 , the node a_1 is not a neighbor of $\langle x_2, x_1, a_2, x_4 \rangle$ and $\langle a_1, S_1 \rangle$ is a Lannér diagram. Hence, $[a_1, x_3] \neq 2$ (a_1 is a neighbor of $S_0 = \langle x_1, x_2, x_3, x_4 \rangle$ and non-neighbor of $\langle x_1, x_2, x_4 \rangle$). The signature condition applied to the diagram $\langle S_0, a_1, a_2, S_1 \rangle$ implies that $[x_3, a_1] = \infty$.

Suppose that a_3 is not a good neighbor of S_1 . Then the dotted edge $\tilde{x}_4 \tilde{a}_3 \subset \Sigma_{S_1}$ remains a dotted edge $x_4 a_3 \subset \bar{S}_1$ in Σ , and the diagram $\langle x_2, x_3, x_4, a_2 \rangle$ of the type A_4 has 4 bad neighbors, x_1, a_1, a_3, y_4 (we use that $[x_3, a_1] = \infty$ and $[a_2, y_4] = \infty$), which is impossible.

Suppose that a_3 is a good neighbor of S_1 . If the edge $x_4 a_3 \subset \bar{S}_1$ is dotted or multiple one, then a_3 remains a bad neighbor of $\langle x_2, x_3, x_4, a_2 \rangle$ and this diagram still has 4 bad neighbors. Therefore, we may assume that $x_4 a_3$ is a simple edge. Hence, S_1 is a diagram of the type H_3 (otherwise $\langle a_2, x_3, x_4, a_3, S_1 \rangle$ is a parabolic diagram of the type $\tilde{B}_6$). So, we have the following diagram:



Furthermore, $[y_4, a_3] \neq \infty$ by Lemma 6.2. If $[y_4, a_3] = 5$ then $\langle y_4, a_3, x_4, x_3 \rangle$ is a diagram of the type H_4 with 4 bad neighbors, a_1, a_2, y_1, x_2 . If $[y_4, a_3] = 4$ then $\langle y_4, a_3, x_4, x_3, x_2 \rangle$ is a diagram of the type B_5 with 4 bad neighbors, a_1, a_2, y_1, x_1 . Hence, $[y_4, a_3] = 2$ or 3 . Similarly, $[a_1, a_3] = 2$ or 3 .

Consider the diagram $\langle S_0, a_3, S_1 \rangle$. The node y_4 is not joined with S_0 . Since the diagram $\langle x_2, x_3, x_4, a_3, y_1, y_2 \rangle$ has 3 bad neighbors (x_1, a_1, y_3), y_4 may be joined with a_3 by a simple edge only. Recall also that $\langle S_1, y_4 \rangle = \bar{S}_0$ is a Lannér diagram, so by Lemma 1.4 we obtain that $[y_4, a_3] \neq 2$. Therefore, $[y_4, a_3] = 3$. It turns out that for any Lannér diagram $\langle y_4, S_1 \rangle$ (where S_1 is of the type H_3) the diagram $\langle S_0, a_3, y_4, S_1 \rangle$ does not satisfy the signature condition unless $[y_4, y_3] = 3$ and $[y_4, y_1] = [y_4, y_2] = 2$. By the similar consideration of the diagram $\langle x_2, x_1, a_2, x_4, a_3, a_1, S_1 \rangle$ we obtain $[a_1, a_3] = [a_1, y_3] = 3$ and $[a_1, y_1] = [a_1, y_2] = 2$. However, in this case the diagram $\langle a_3, y_4, y_3, y_2 \rangle$ of the type H_4 has 4 bad neighbors (a_2, x_4, y_1, a_1).

Case 2.2.4. Suppose that Σ_{S_1} is one of the diagrams shown in Fig. 14(e)-(g). In these cases a_2 still cannot be a good neighbor of S_1 , therefore, it is not a neighbor and $[a_2, y_4] = \infty$ as it was shown above. At the same time, if $[x_3, a_3] = 2$, then a_3 may be a good neighbor of S_1 and it may occur that $[x_4, a_3] \neq \infty$ in Σ .

Suppose that $[x_4, a_3] \neq \infty$ in Σ , and hence, $[x_3, a_3] = 2$. Consider the subdiagram $Y = \langle S_0, a_2, a_3, S_1 \rangle$. If $[x_4, a_3] = 3$ or 5 , then Y does not satisfy the signature condition (namely, at least one of the subdiagrams $Y \setminus a_2$ and $Y \setminus y_4$ does not satisfy the condition). If $[x_4, a_3] = 4$ then Y contains a parabolic subdiagram $\langle x_2, x_3, x_4, a_3, y_1 \rangle$ of the type $\tilde{F}_4$ (here we assume that the nodes of S_1 are numbered in such a way that $[y_1, y_2] = 3$, $[y_2, y_3] = 4$ or 5).

Therefore, $[x_4, a_3] = \infty$. Consider the subdiagram $S_7 = \langle x_2, x_1, a_2 \rangle$ of the type H_3 . It has at least 2 bad neighbors x_3 and y_4 . Furthermore, the dotted edge $x_4 a_3$ is not joined with S_7 , so $\langle x_4, a_3 \rangle \subset \bar{S}_7$, which implies that S_7 has no extra bad neighbors. In particular, a_1 is not a bad neighbor of S_7 . Suppose that a_1 is a good neighbor of S_7 . Then $S_8 = \langle a_1, S_7 \rangle$ is a diagram of the type H_4 with 3 bad neighbors (x_3, y_4 and some node of S_1 joined with a_1). However, $\bar{S}_8$ contains a dotted edge $x_4 a_3$. The contradiction shows that a_1 is not a neighbor of S_7 . Thus, the diagram $\bar{S}_7 = \Sigma_{S_7} = \langle x_4, a_3, a_1, S_1 \rangle$ is a diagram of a 4-prism, which implies that $\langle a_1, S_1 \rangle$ is a Lannér diagram. A Lannér diagram $\langle a_2, x_1, x_2, x_3 \rangle$ should be joined with a Lannér diagram $\langle a_1, S_1 \rangle$. Since a_1 is not a neighbor of S_7 , this implies that $[a_1, x_3] \neq 2$. Consider the

subdiagram $S_9 = \langle a_2, x_2, x_3, x_4 \rangle$ (of the type $A_1 + A_3$ or A_4 or D_4 in cases (e), (f) and (g) respectively). It has 3 bad neighbors x_1, a_3, y_4 , so a_1 is a good neighbor of S_9 , which implies $[a_1, x_3] = 3$, $[a_1, x_4] = 2$.

Finally, consider the diagram $Z = \langle S_0, a_2, a_1, S_1 \rangle$. This diagram contains no dotted edges, so we have finitely many possibilities for Z . Moreover, Z satisfies the following conditions: $\langle S_0, S_1 \rangle$ is a diagram of the type $H_4 + H_3$ or $H_4 + B_3$; a_2 is not joined with S_1 and is joined with S_0 in one of the ways shown in Fig. 14(e)-(g); $[a_1, x_3] = 3$ and $a_1 x_3$ is a unique edge connecting a_1 with $\langle a_2, S_0 \rangle$; $\langle a_1, S_1 \rangle$ is a Lannér diagram. However, no diagram satisfies these conditions together with the signature condition.

Case 3. Suppose that S_1 has 3 bad neighbors. Let $S'_1 \subset \overline{S}_0$ be a subdiagram of the type H_3 or B_3 , $S'_1 \neq S_1$ (see Lemma 1.5). We denote $\langle y_1, y_2, y_3 \rangle = S_1$, $\langle y_2, y_3, y_4 \rangle = S'_1$. By Cases 1 and 2 we may assume that both S_1 and S'_1 have 3 bad neighbors. We have two possibilities (up to permutation of the nodes a_1, a_2, a_3): either a_1 and a_2 are bad neighbors of S_1 as well as of S'_1 (in addition to y_4 and y_1 respectively), or y_4, a_1, a_2 are bad neighbors of S_1 and y_1, a_2, a_3 are bad neighbors of S'_1 .

Suppose that a_1 and a_2 are bad neighbors for each of S_1 and S'_1 . Then the node a_3 is not a bad neighbor for both S_1 and S'_1 . The diagram Σ_{S_1} is a Lannér diagram of order 5. By Cor. 1.1, $\overline{S}_1 = \langle S_0, a_3 \rangle$ is a Lannér diagram, too. So, the diagram $\langle S_0, a_3, \overline{S}_0 \rangle$ consists of a Lannér diagram $\langle S_0, a_3 \rangle$ (where S_0 is of the type H_4), and a Lannér diagram $\overline{S}_0$, the node a_3 is not a bad neighbor for both S_1 and S'_1 . The only such a diagram satisfying the signature condition is the diagram shown in Fig. 12(b). However, by Proposition 1.11, the edge $\tilde{x}_4 \tilde{a}_3$ of Σ_{S_1} is a dotted edge, which contradicts the fact that Σ_{S_1} is a Lannér diagram of order 5.

Therefore, we may assume that y_4, a_1, a_2 are bad neighbors of S_1 , and y_1, a_2, a_3 are the bad neighbors of S'_1 . By consideration of $\overline{S}_1$ and $\overline{S}'_1$ we conclude that $\langle S_0, a_1 \rangle$ and $\langle S_0, a_3 \rangle$ are Lannér diagrams.

Now, we consider two cases: either both a_1 and a_3 are joined with $\overline{S}_0$ by dotted edges, or at least one of a_1 and a_3 (say a_1) is joined with $\overline{S}_0$ by non-dotted edges only.

Case 3.1. Suppose that both a_1 and a_3 are joined with $\overline{S}_0$ by dotted edges. Since a_1 is a bad neighbor of S_1 and it is not a bad neighbor of S'_1 , the dotted edge joining a_1 with $\overline{S}_0$ is $a_1 y_1$. Similarly, $[a_3, y_4] = \infty$. By Lemma 6.2, $[a_1, a_3] \neq \infty$. Hence, we have finitely many possibilities for the diagram $\langle S_0, a_1, a_3 \rangle$ (4 possibilities for each of Lannér diagrams $\langle S_0, a_1 \rangle$ and $\langle S_0, a_3 \rangle$ and 3 possibilities for $[a_1, a_3] \in \{3, 4, 5\}$).

Consider the subdiagram $S_{10} = \langle x_2, x_3, x_4, a_1 \rangle$ of the type D_4, A_4, B_4 or H_4 . If a_3 is a bad neighbor of S_{10} then S_{10} has 3 bad neighbors x_1, y_1 and a_3 , thus, a_2 is not a bad neighbor of S_{10} . If a_3 is not a bad neighbor of S_{10} , then the diagram $\langle S_{10}, a_3 \rangle$ has 3 bad neighbors x_1, y_1, y_4 , and a_2 is not a bad neighbor of $\langle S_{10}, a_3 \rangle$. In any case, a_2 is not a bad neighbor of S_{10} . Similarly, a_2 is not a bad neighbor of $S_{11} = \langle x_2, x_3, x_4, a_3 \rangle$.

Suppose that a_2 is a bad neighbor of the diagram $S_{12} = \langle x_1, x_2, x_3 \rangle$ of the type H_3 . Then $P(S_{12})$ is a 4-polytope with at most $4 + 3$ facets. However, $\Sigma_{S_{12}} \neq \overline{S}_{12}$, and $\Sigma_{S_{12}}$ contains 4 dotted edges $\tilde{y}_4 \tilde{a}_3, \tilde{a}_3 \tilde{x}_4, \tilde{x}_4 \tilde{a}_1, \tilde{a}_1 \tilde{y}_1$, which is impossible for a diagram of a 4-polytope with at most 7 facets. Therefore, a_2 is not a bad neighbor of S_{12} , which implies that a_2 is not a neighbor of $x_1 x_2$. Hence, a_2 is a neighbor of $x_3 x_4$. Thus, each of S_3 and S_{11} is a diagram of the type D_4 or A_4 .

Suppose that S_{10} and S_{11} are of the same type. Then $[a_1, a_3] = 2$ (otherwise either $[a_1, a_3] = \infty$ in contradiction to Lemma 6.2, or $\langle a_1, a_3 \rangle$ has 3 bad neighbors in contradiction to Lemma 6.1, or one of the diagrams $\langle x_3, a_1, a_3 \rangle$ and $\langle x_4, a_1, a_3 \rangle$ is a parabolic diagram of the type $\tilde{A}_2$). This implies that $\langle S_{10}, a_3, a_2 \rangle$ contains a parabolic subdiagram. Hence, S_{10} and S_{11} are of different types. If $[a_1, a_3] = 2$ or 3 then $\langle S_{10}, a_3, a_2 \rangle$ still contains a parabolic diagram. If $[a_1, a_3] = 4$ or 5 then the diagram $S_{13} = \langle a_1, a_3 \rangle$ has 2 bad neighbors, so $\Sigma_{S_{13}}$ is a diagram of a 5-prism.

However, in $\Sigma_{S_{13}}$ the subdiagram $\langle \tilde{x}_1, \tilde{x}_2, \tilde{x}_3, \tilde{x}_4 \rangle$ is a linear diagram in which one triple edge and one simple edge are joined by the dotted edge $\tilde{x}_2 \tilde{x}_3$. No diagram of a 5-prism contains such a subdiagram, which shows that the case 3.1 is impossible.

Case 3.2. Suppose that a_1 is joined with S_0 by non-dotted edges only. Then the subdiagram $T = \langle S_0, a_1, \bar{S}_0 \rangle$ contains no dotted edges, so we have finitely many possibilities for the diagram T . Notice also that T satisfies the following properties:

- S_0 is a diagram of the type H_4 , and $\langle S_0, a_1 \rangle$ is a Lannér diagram;
- $\bar{S}_0$ is a Lannér diagram of order 4, $\bar{S}_0$ is not joined with S_0 ;
- S_1 and S'_1 are subdiagrams of $\bar{S}_0$ or the types H_3 or B_3 ;
- a_1 is a bad neighbor of S_1 and not a bad neighbor of S'_1 .

However, there are only two diagrams satisfying these properties together with signature condition, namely the diagrams shown in Fig. 15(a) and 15(b).

Consider the diagram shown in Fig. 15(a). The subdiagram $\langle x_2, x_3, x_4, a_1, y_1 \rangle$ of the type B_5 has 3 bad neighbors in Σ : x_1, y_2 and a_3 . Hence, a_2 is not a bad neighbor of $\langle x_2, x_3, x_4, a_1, y_4 \rangle$, which implies that a_2 is joined with $\langle x_1, x_2 \rangle$. Therefore, a_2 is a bad neighbor of $S_{14} = \langle x_1, x_2, x_3 \rangle$, and $P(S_{14})$ is a 4-polytope with $4 + 3$ facets (it is easy to see that S_{14} has no other bad neighbors). However, $\Sigma_{S_{14}}$ contains $\langle \tilde{a}_1, \tilde{y}_1, \tilde{y}_2 \rangle$, the subdiagram composed of a double edge $\tilde{a}_1 \tilde{y}_1$ adjacent to a triple edge $\tilde{y}_1 \tilde{y}_2$, which is impossible for a diagram of a 4-polytope with $4 + 3$ facets.

In particular, we obtain that for any diagram $S \subset \Sigma$ of the type H_4 the diagram $Q = \langle S, \bar{S}, a_1 \rangle$ is the diagram shown in Fig. 15(b) and $\bar{S}$ is a linear Lannér diagram with one double, one simple and one triple edge.




Figure 15:

Consider the diagram $\langle Q, a_3 \rangle$. It contains at most 1 dotted edge (Lemma 6.2), which cannot have endpoint neither in S_0 ($\langle S_0, a_3 \rangle$ is a Lannér diagram) nor in S_1 (a_3 is not a bad neighbor of S_1). So, either $\langle S_0, a_3, y_4, S_1 \rangle$ or $\langle S_0, a_3, a_1, S_1 \rangle$ is a diagram containing no dotted edges. A direct check shows that the former diagram never satisfies signature condition, while the latter satisfies it in a unique case shown in Fig. 15(c). Moreover, $[a_3, y_4] = \infty$, otherwise $\det(\langle S_0, a_1, a_3, \bar{S}_0 \rangle) \neq 0$.

Now, we are left to determine how it is possible to attach a_2 to the diagram $\langle S_0, a_1, a_3, \bar{S}_0 \rangle$. Consider the subdiagram $S_{15} = \langle y_1, y_2, a_1, x_4 \rangle$ of the type H_4 with its bad neighbors x_3, a_3, y_2 . As it was shown above, the diagram $\bar{S}_{15} = \langle x_1, x_2, a_2, y_4 \rangle$ is a linear Lannér diagram with one double, one simple and one triple edge. Hence, either $[a_2, x_2] = 2, [a_2, x_1] = 3, [a_2, y_4] = 4$ or $[a_2, x_2] = 3, [a_2, x_1] = 2, [a_2, y_4] = 4$. Furthermore, a_2 is not joined with $\langle x_3, x_4, a_3, a_1 \rangle$ since the diagram $\langle x_2, x_3, x_4, a_3, a_1, y_2 \rangle$ of the type E_6 already has 3 bad neighbors x_1, y_1, y_4 . Also, a_2 is not joined with $\langle y_1, y_2 \rangle$ since the diagram $\langle y_1, y_2, a_1, x_4 \rangle$ of the type H_4 has 3 bad neighbors x_3, a_3, y_3 . Since a_2 is a bad neighbor of S_1 , $[a_2, y_3] \neq 2$. If $[a_2, y_3] = 3$, then $S_{16} = \langle x_2, x_1, a_2, y_3 \rangle$ is a diagram of the type H_4 such that $\bar{S}_{16} = \langle y_1, a_1, x_4, a_3 \rangle$ is a non-connected diagram, which is impossible. If $[a_2, y_3] = 5$ (or 4) then $\langle a_2, y_3, y_2, a_1 \rangle$ (or $\langle a_2, y_3, y_2, a_1, x_4, a_3 \rangle$) is a diagram of the type H_4 (B_6) with 4 bad neighbors y_4, y_1, x_4 and one of x_1 and x_2 . Hence, $[a_2, y_3] = \infty$.

Therefore, a_2 is an end of three edges: one dotted edge a_2y_3 , one double edge a_2y_4 and one simple edge (either a_2x_1 or a_2x_2). If $[a_2, x_1] = 3$ then the subdiagram $\langle S_0, a_1, a_2, a_3, y_1, y_2 \rangle$ does not satisfy the signature condition (it contains no dotted edges, so we check the signature directly). Thus, $[a_2, x_1] = 2$, $[a_2, x_2] = 3$ and we arrive with the diagram Σ_{P_7} .

All cases are considered, the only polytope P_7 is found. So, the lemma is proved. □

References

- [An] E. Andreev, *Intersection of plane boundaries of acute-angled polyhedra*, Math. Notes 8 (1971), 761–764.
- [Al] D. Allcock, *Infinitely many hyperbolic Coxeter groups through dimension 19*, Geometry & Topology 10 (2006), 737–758.
- [Bo] R. Borcherds, *Coxeter groups, Lorentzian lattices, and K3 surfaces*, IMRN 19 (1998), 1011–1031.
- [Bu1] V. O. Bugaenko, *Groups of automorphisms of unimodular hyperbolic quadratic forms over the ring $\mathbb{Z}\left[\frac{\sqrt{5}+1}{2}\right]$* , Moscow Univ. Math. Bull. 39 (1984), 6–14.
- [Bu2] V. O. Bugaenko, *Arithmetic crystallographic groups generated by reflections, and reflective hyperbolic lattices*, Adv. Sov. Math. 8 (1992), 33–55.
- [C] H. S. M. Coxeter, *Discrete groups generated by reflections*, Ann. Math. 35 (1934), 588–621.
- [E1] F. Esselmann, *Über kompakte hyperbolische Coxeter-Polytope mit wenigen Facetten*, Universität Bielefeld, SFB 343, Preprint 94-087.
- [E2] F. Esselmann, *The classification of compact hyperbolic Coxeter d -polytopes with $d + 2$ facets*, Comment. Math. Helvetici 71 (1996), 229–242.
- [FT1] A. Felikson, P. Tumarkin, *Reflection subgroups of reflection groups*, Funct. An. Appl. 38 (2004), 313–314.
- [FT2] A. Felikson, P. Tumarkin, *On hyperbolic Coxeter polytopes with mutually intersecting facets*, arXiv:math.MG/0604248.
- [G] B. Grünbaum, *Convex polytopes*, John Wiley & Sons, 1967.
- [ImH] H.-C. Im Hof, *Napier cycles and hyperbolic Coxeter groups*, Bull. Soc. Math. de Belg. Série A, XLII (1990), 523–545.
- [K] I. M. Kaplinskaya, *Discrete groups generated by reflections in the faces of simplicial prisms in Lobachevskian spaces*, Math. Notes 15 (1974), 88–91.
- [L] F. Lannér, *On complexes with transitive groups of automorphisms*, Comm. Sem. Math. Univ. Lund 11 (1950), 1–71.
- [M] V. S. Makarov, *On Fedorov groups of four- and five-dimensional Lobachevsky spaces*, Studies in General Algebra, no. 1, Kishinev St. Univ., 1970, 120–129 (Russian).
- [P] M. N. Prokhorov, *The absence of discrete reflection groups with non-compact fundamental polyhedron of finite volume in Lobachevsky space of large dimension*, Math. USSR Izv. 28 (1987), 401–411.

- [R] O. P. Rusmanov, *Examples of non-arithmetic crystallographic Coxeter groups in n -dimensional Lobachevsky space for $6 \leq n \leq 10$* , Problems in Group Theory and Homology Algebra, Yaroslavl, 1989, 138–142 (Russian).
- [T] P. Tumarkin, *Compact hyperbolic Coxeter n -polytopes with $n + 3$ facets*, arXiv:math.MG/0406226.
- [V1] E. B. Vinberg, *On groups of unit elements of certain quadratic forms*, Math. USSR Sb. 16 (1972), 17–35.
- [V2] E. B. Vinberg, *The absence of crystallographic groups of reflections in Lobachevsky spaces of large dimension*, Trans. Moscow Math. Soc. 47 (1985), 75–112.
- [V3] E. B. Vinberg, *Hyperbolic reflection groups*, Russian Math. Surveys 40 (1985), 31–75.
- [V4] E. B. Vinberg (Ed.), *Geometry II*, Encyclopedia of Mathematical Sciences, Vol. 29, Springer-Verlag Berlin Heidelberg, 1993.
- [VK] E. B. Vinberg, I. M. Kaplinskaya, *On the groups $O_{18,1}(\mathbb{Z})$ and $O_{19,1}(\mathbb{Z})$* , Soviet Math. Dokl. 19 (1978), 194–197.

Independent University of Moscow

e-mail: felikson@mccme.ru

pasha@mccme.ru